

VU LEARNING HUB

vulearninghub.xyz

HIGHLIGHTED HANDOUT

BIO504T

BIO504T

BIO504T by VU Learning HUB

Prepared for VU students • Free study resources

<https://vulearninghub.xyz>



Virtual University of Pakistan

FEDERAL GOVERNMENT UNIVERSITY

COURSE HANDOUTS

Biochemistry-I

BIO504T

Order Your VU Books Now

Get all Virtual University books in high-quality hard copy, delivered right to your doorstep. Visit vubookshop.pk



Order Now

BIO504T Biochemistry-I

Topic no 1 to 112 in mid term and 113 to 225 final term syllabus

Lesson 1 Carbohydrates

Carbohydrates are polyhydroxy aldehydes or ketones, or substances that yield such compounds on hydrolysis. Many, but not all, carbohydrates have the empirical formula $(CH_2O)_n$; some also contain nitrogen, phosphorus or sulfur.

Classification

"Carbohydrates are classified into **four** major groups:

1. Monosaccharides
2. Oligosaccharides
3. Disaccharides
4. Polysaccharides

Monosaccharides (simple sugars) are those which cannot be hydrolyzed further into simpler forms. The backbones of common monosaccharides are *unbranched* carbon chains in which all the carbon atoms are linked by single bonds. In **the open-chain form**, one of the carbon atoms is double-bonded to an oxygen atom to form a **carbonyl group**. Each of the other carbon atoms has a hydroxyl group.

If the carbonyl group is at an end of the carbon chain (that is, in an aldehyde group) the monosaccharide is an **aldose**.

If the carbonyl group is at any other position (in a ketone group) the monosaccharide is a **ketose**. The **simplest** monosaccharides are the two three-carbon trioses:

Glyceraldehyde, an aldotriose

Dihydroxyacetone, a ketotriose

The carbons of a sugar are numbered beginning at the end of the chain nearest the carbonyl group.

All common

wxyx•kmm k°no•"kxn"nə kmm k°no•"r k\o"kwō•"oxnsxq"Δsr "r o"•#ps" -y•o3""

Ketoses are designated by inserting "ul" into the name of a corresponding aldose.

For example, ribulose is the ketopentose corresponding to the aldopentose ribose. "

There are a few exceptions however, for example, fructose and dihydroxyacetone.

Monosaccharides can be subdivided further depending upon:

Number of carbon atoms. Whether aldehyde or ketone groups are present as *aldoses* or *ketoses*.

General formula	Aldosugars	Ketosugars
Trioses $C_3H_6O_3$	Glyceraldehyde	Dihydroxyacetone
Tetroses $C_4H_8O_4$	Erythrose	Erythrulose
Pentoses $C_5H_{10}O_5$	Ribose	Ribulose
Hexoses $C_6H_{12}O_6$	Glucose	Fructose

The **hexoses** which include the aldohexose D-glucose and the ketohexose D-fructose, are the most common monosaccharides in nature. The **aldopentoses** D-ribose and 2-deoxy-D-ribose are components of nucleotides and nucleic acids.

Lesson 2

Disaccharides are those sugars which yield two molecules of the same or different monosaccharides on hydrolysis.

Maltose - 2 molecules of glucose on hydrolysis

Lactose 1 glucose & 1 galactose on hydrolysis

Sucrose 1 glucose & 1 fructose on hydrolysis

Oligosaccharides are short chains of monosaccharide units or residues (3-10) joined by characteristic linkages called glycosidic linkage. **Glycosidic bond** of maltose is formed between the OH of carbon 1 and carbon 4 of 2nd glucose monomers. Therefore, it forms an alpha (1 → 4) glycosidic bond.

Lesson 3

Polysaccharides (Glycans) are those which yield more than 10 molecules of monosaccharides on hydrolysis. Some have hundreds or thousands of units. Some polysaccharides, such as **cellulose**, are linear chains; others, such as **glycogen**, are branched.

Both glycogen and cellulose consist of recurring units of D-glucose, but they differ in the type of glycosidic linkage and consequently have strikingly different properties and biological roles.

Polysaccharides are further divided into two groups:

1. Homopolysaccharides (homoglycans): Polymers of same monosaccharide units e.g. starch, glycogen, inulin, cellulose, dextrans, dextrans

2. Heteropolysaccharides (heteroglycans): Polymer of different monosaccharide units or their derivatives e.g. Mucopolysaccharides (glycosaminoglycans)

Biomedical Importance

Chief source of energy. Constituents of compound lipids and conjugated proteins. Lactose, the principal sugar of milk. Degradation products utilized for synthesis of other substances such as fatty acids, amino acids, cholesterol, etc. Constituents of mucopolysaccharides which form the ground substance of mesenchymal tissues. Certain carbohydrate derivatives are used as drugs like **cardiac glycosides**. Inherited deficiency of certain enzymes in metabolic pathways of different carbohydrates can cause diseases e.g. **galactosemia, glycogen storage diseases, lactose intolerance**, etc.

Derangement of glucose metabolism is seen in **Diabetes mellitus**.

Lesson 4

Isomerism

"The existence of two or more compounds having the same molecular formula but a different arrangement of atoms within the molecule."

Isomerism

Compounds which are identical in composition and differ only in configuration (the fixed spatial arrangement of atoms) are called **stereoisomers**

Stereoisomers that are mirror images of each other are called **enantiomers**

Pairs of stereoisomers that are not mirror images of each other are called **diastereomers**

The identifying characteristic of stereoisomers is that they cannot be interconverted without temporarily breaking one or more covalent bonds

A carbon atom to which four different atoms or groups of atoms are attached is said to be **asymmetric** or **chiral**

Lesson 5

Configuration is conferred by the presence of either

(a) **chiral centers**, around which substituent groups are arranged in a specific orientation

(b) **double bonds**, around which there is no freedom of rotation

"Geometric isomers; differ in the arrangement of their substituent groups with respect to the nonrotating double bond (each is a well-defined compound that can be separated from the other, and each has *its own unique chemical properties*)" The identifying characteristic of configurational isomers is that they cannot be interconverted without temporarily breaking one or more covalent bonds.

The configurations of maleic acid and its isomer, fumaric acid.

These compounds are **geometric, or cis-trans, isomers;**

They differ in the arrangement of their substituent groups with respect to the nonrotating double bond

Lesson 6

D and L isomerism:

The designation of a sugar isomer as the D form or as the L form is determined by its spatial relationship to the parent compound of the carbohydrates, the glyceraldehyde. The orientation of the H and OH groups around the **carbon atom adjacent to the terminal primary alcohol or carbon most distant from the carbonyl carbon the penultimate carbon**

(carbon 5 in glucose) determines whether the sugar belongs to the D or L series.

When the OH group on this carbon is on the right the sugar is the D isomer;

when it is on the left, it is the L isomer. Of the 16 possible aldohexoses eight are D forms and eight are L. Most of the hexoses of living organisms are D isomers and the enzymes responsible for their metabolism are specific for this configuration.

Lesson 7

Four different substituents bonded to a chiral carbon atom may be arranged in two different ways in space, that is, have two stereoisomers (with similar or identical chemical properties **but differing in certain physical and biological properties**).

A molecule with only one chiral carbon can have two stereoisomers; when two or more (n) chiral carbons are present, there can be 2^n stereoisomers.

Enantiomers have nearly identical chemical properties but differ in a characteristic physical property:

- their interaction with plane-polarized light

Thus the presence of asymmetric carbon atoms also confers **optical activity** on the compound.

When a beam of plane-polarized light is passed through a solution of an **optical isomer**, it rotates either to the right, dextrorotatory (+), or to the left, levorotatory (−). The direction of rotation of plane polarized light is independent of the stereochemistry of the sugar, so it may be designated D(−), D(+), L(−), or L(+).

For example, the naturally occurring form of fructose is the D(−) isomer.

In solution, glucose is dextrorotatory, and glucose solutions are sometimes known as **dextrose**.

Lesson 8

When equal amounts of dextrorotatory and levorotatory isomers are present, the resulting mixture has no optical activity

Such a mixture is said to be **racemic** and separation of optically active isomers from a racemic mixture is called **resolution**

All the monosaccharides except dihydroxyacetone contain one or more asymmetric

(chiral) carbon atoms and thus occur in optically active isomeric forms

The simplest aldose, glyceraldehyde contains one chiral center (the middle carbon atom) and therefore has two different optical isomers

Lesson 9

In aqueous solution, aldotetroses and all monosaccharides with five or more carbon atoms in the backbone occur predominantly as **cyclic (ring) structures**.

The formation of these ring structures is the result of a general reaction between alcohols and aldehydes or ketones to form derivatives called **hemiacetals or hemiketals**.

Which contain an additional **asymmetric carbon atom** and thus can exist in two stereoisomeric forms

For example, D-glucose exists in solution as an **intramolecular hemiacetal** in which the free hydroxyl group at C-5 has reacted with the aldehydic C-1, therefore rendering the latter carbon (aldehydic C-1) asymmetric. And producing two stereoisomers, designated **"α -"**

Carbon 1, after cyclization has four different groups attached to it and thus it also becomes asymmetric

Isomeric forms of monosaccharides that differ only in their configuration about the hemiacetal or hemiketal carbon atom are called **anomers**

The carbonyl carbon atom is called the **anomeric carbon**

Lesson 10

Ring Structure of Carbohydrates

For example, D-glucose exists in solution as an **intramolecular hemiacetal** in which the free hydroxyl group at C-5 has reacted with the aldehydic C-1, "Therefore rendering the latter carbon (aldehydic C-1) asymmetric

And producing two stereoisomers, designated **" α -"**

Carbon 1, after cyclization has four different groups attached to it and thus it also

Isomeric forms of monosaccharides that differ only in their configuration about the hemiacetal or hemiketal carbon atom are called **anomers**

Lesson 11

The ring structures of monosaccharides are similar to the ring structures of either pyran (six-membered ring) and are called **pyranoses** or furan (a five-membered ring), which are known as **furanoses**

The six-membered aldopyranose ring is much more stable than the aldofuranose ring
Therefore predominates in aldohexose and aldopentose solutions

Similarly Crystalline glucose is **"D -glucopyranose**

[α and β anomers of D-glucose interconvert in aqueous solution by a process called **mutarotation**.

Thus, a solution of α -D-glucose and a solution of β -D-glucose eventually form identical equilibrium mixtures having identical optical properties.

This mixture consists of about:

one-third α -D-glucose, two-thirds β -D-glucose, and very small amounts of the linear and five-membered ring (glucofuranose) forms.

"kxn" "p°w•"k°o"UV["ws°°y°°"swkqo•"kxn"k°o"°o°°on"y°k•"diastereomers

Enzymes are able to distinguish between these two structures and use one or the other preferentially

My "o-kwzwlqvnyqox"s "x'ro\$on"pyw" -D-glucopyranose, whereas cellulose is "x'ro\$on"pyw" -D-glucopyranose

Ro'yr o-y•o• "kv y"ymf°"x" "kxn" "kxywo°sm"p°w•

In these compounds the hydroxyl group at C-5 (or C-6) reacts with the keto group at C-2, forming a furanose (or pyranose) ring containing a hemiketal linkage

D-Fructose readily forms the furanose ring ; the more common anomer of this sugar in combined forms or in derivatives is β -D-fructofuranose

Lesson 12 Epimers

Two sugars which differ from one another only in configuration around a single carbon atom are termed **epimers**

Glucose and Galactose are an example of an epimeric pair which differ only with respect to C_4

Similarly Glucose and mannose are epimers with respect to C_2

However, Galactose and mannose are NOT epimers as they differ in the position of OH groups at two carbons (2 and 4), and are therefore defined only as isomers

Trioses:

Both **D-Glyceraldehyde & Dihydroxyacetone** occur as intermediates in glycolysis
They are also the precursors of glycerol

Tetroses:

Erythrose occurs as an intermediate in the Hexose Mono Phosphate shunt which is an alternative pathway for glucose oxidation

Pentoses:

D-Ribose is a constituent of nucleic acid *RNA* and also certain co-enzymes like *FAD*, *NAD* & *coenzyme A*

D-2-Deoxyribose is a constituent of *DNA*

Hexoses

D-Glucose (Dextrose, Grape sugar):

Most dietary carbohydrate is absorbed into the bloodstream as glucose formed by hydrolysis of dietary starch and disaccharides, and other sugars are converted to glucose in the liver

Sources include fruit juices, hydrolysis of starch, cane or beet sugar, maltose and lactose

Glucose is the major metabolic fuel of mammals

It is the chief physiological sugar present in normal blood at a fairly constant level

All tissues utilize glucose for energy

Erythrocyte utilize glucose solely for energy purposes and Brain cells rely heavily on glucose

It is the precursor for synthesis of all the other carbohydrates in the body, including **glycogen** for storage; **ribose** and **deoxyribose** in nucleic acids; **galactose** in lactose of milk, in glycolipids, And in combination with protein in glycoproteins and proteoglycans

Stored as Glycogen in liver and muscles only Excreted in the urine (glucosuria) in poorly controlled diabetes mellitus as a result of hyperglycemia

Lesson 13 D-Galactose:

Occurs as a constituent of milk sugar lactose

It is an epimer of glucose and differs in orientation of H and OH around C-4

Readily metabolized to glucose;

Synthesized in the mammary gland for synthesis of lactose in milk

A constituent of glycolipids and glycoproteins

Hereditary galactosemia as a result of failure to metabolize galactose leads to cataracts

D-Fructose:

Commonly known as *fruit sugar* as it occurs free in fruits, honey etc

Also occurs as a constituent of sucrose

It is levorotatory and hence is also called Levulose

Seminal fluid is rich in fructose and sperm utilize fructose for energy

Readily metabolized either via glucose or directly

Hereditary fructose intolerance leads to fructose accumulation

Properties of Monosaccharides

1. Oxidation of the carbonyl (aldehyde) carbon of glucose to the carboxyl level produces **gluconic acid** " Other aldoses yield other **aldonic acids**

Oxidation of the carbon at the other end of the carbon chain---C-6 of glucose, galactose, or mannose forms the corresponding **uronic acid**, glucuronic, galacturonic, or mannuronic acid

The carboxylic acid groups of the acidic sugar derivatives are ionized at pH 7, and the compounds are therefore correctly named as the carboxylates-glucuronate, galacturonate, and so forth

Both aldonic and uronic acids form stable intramolecular esters called **lactones**

D-Glucono- δ -lactone results from formation of an ester linkage between the C-1 carboxylate group and the C-5 hydroxyl group of gluconate

Glucuronic acid(Oxidation at C-1)

Gluconic acid(Oxidation at C-1)

Lesson 14 to 21

Properties of Monosaccharides

Glucono delta-lactone (GDL) is commonly found in honey, fruit juices, personal lubricants, and wine. GDL is neutral.

But hydrolyses in water to gluconic acid which is acidic, adding a tangy taste to foods

It is metabolized to glucose; one gram of GDL yields roughly the same amount of metabolic energy as one gram of sugar.

In the body, D-Glucuronic acid is formed from glucose in liver

It occurs as a constituent of certain polysaccharides

It is of importance in that it conjugates toxic substances, drugs, hormones and even bilirubin.

And converts them to a soluble non toxic substance, a glucuronide, which is excreted in urine " Oxidation of the carbon at the both ends of the carbon chain.

The terminal methyl & the carbonyl carbon of glucose, galactose, or mannose-forms the corresponding

aldaric acid

Glucose forms **Glucaric acid**

Monosaccharides can be oxidized by relatively mild oxidizing agents such as cupric (Cu^{2+}) ion

The carbonyl carbon is oxidized to a carboxyl group

Glucose and other sugars capable of reducing cupric ion are called

reducing sugars **Sugars as reducing agents**

Oxidation of the anomeric carbon (and probably the neighboring carbon) of glucose and other sugars under alkaline conditions is the basis for Fehling's reaction

Benedict's reaction

The cuprous ion (Cu^+) produced forms a red cuprous oxide precipitate

By measuring the amount of oxidizing agent reduced by a solution of a sugar, it is also possible to estimate the concentration of that sugar

For many years this test was used to detect and measure elevated glucose levels in blood and urine in the diagnosis of diabetes mellitus

Properties of Monosaccharides

2. Condensation of phosphoric acid with one of the hydroxyl groups of a sugar forms a **phosphate ester**, as in glucose 6-phosphate

Sugar phosphates are relatively stable at neutral pH and bear a negative charge

In fact, one of the strategies of glycolysis is to form three carbon intermediates, that can transfer their phosphate groups to ADP, to achieve a net synthesis of ATP

In the synthesis and metabolism of carbohydrates, the intermediates are very often not the sugars themselves but their phosphorylated derivatives

Another function of phosphorylation is the creation of reactive intermediates for the formation of O- and N- glycosidic linkages

Properties of Monosaccharides

3. Synthesis of Amino Sugar

If the hydroxyl at C-2 of the parent monosaccharide is replaced with an amino group, the corresponding **amino sugar** is formed

The amino sugars include:

D -glucosamine, a constituent of hyaluronic acid

D -galactosamine (also known as *chondrosamine*), a constituent of chondroitin

D -mannosamine

The amino group is nearly always condensed with **acetic acid**, as in N-acetylglucosamine

Both glucosamine and galactosamine occur as N-acetyl derivatives in more complex carbohydrates.

For example glycoproteins and is part of many structural polymers, including those of the bacterial cell wall

Bacterial cell walls also contain a derivative of glucosamine, **N-acetylmuramic acid**, in which lactic acid (a three carbon carboxylic acid) is ether-linked to the oxygen at C-3 of N-acetylglucosamine

The Amino Sugars (Hexosamines) are Components of Glycoproteins, Gangliosides, & Glycosaminoglycans

Several **antibiotics** (e.g, erythromycin) contain amino sugars, which are important for their antibiotic activity

4. When oxygen of a

OH group is removed in a monosaccharide, leaving behind only hydrogen, **deoxy sugars** are formed

2-Deoxy-D-Ribose is derived from D-Ribose and is an important constituent of DNA

The substitution of a hydrogen for the hydroxyl group at C-6 of ;
L-galactose &
L-mannose produces;
L-fucose and
L-rhamnose, respectively.

L-Fucose is found in the complex oligosaccharide components of glycoproteins and glycolipids;

L-Rhamnose is found in plant polysaccharide

5. Glycosides are formed by condensation between the hydroxyl group of the anomeric carbon of a monosaccharide, and a second compound that may or may not (in the case of an **aglycone**) be another monosaccharide. If the second group is a **hydroxyl**, the **O-glycosidic** bond is an **acetal link** because it results from a reaction between a hemiacetal group (formed from an aldehyde and an OH group) and another OH group.

If the hemiacetal portion is glucose, the resulting compound is a **glucoside**. If galactose, a **galactoside**, and so on.

6. The monosaccharides may be reduced to form their corresponding alcohols by reducing agents;

Examples are

- D-Glucose yields D-Sorbitol
- D-Galactose yields D-Dulcitol
- D-Mannose yields D-Mannitol
- Ribose to ribitol

Sorbitol:

35-60% sweetness of glucose

Used as an artificial sweetener

Accumulates in tissues such as the eye lens in diabetes mellitus **Mannitol:**

Administered intravenously as an osmotic diuretic in acute renal failure

Also used to relieve raised intracranial pressure by forced diuresis


7. Tautomerization

The process of shifting a hydrogen atom from one carbon atom to another to produce **enediols** is known as tautomerization

Sugars possessing anomeric carbon atom undergo tautomerization in alkaline solutions

When glucose is kept in alkaline solution for several hours, it undergoes isomerization to form Dfructose and D-mannose

This reaction results in the formation of a common intermediate-namely enediol--for all the three sugars
The enediols are highly reactive, hence sugars in alkaline solution are powerful reducing agents



8. Osazone formation:

It is a useful means of preparing *crystalline derivatives* of sugars called **osazones**

They are obtained by adding a mixture of **phenyl hydrazine hydrochloride** and **sodium acetate** to a sugar solution and heating in water bath for 45 mins

Crystals are formed after the solution is cooled slowly

Reaction involves only the carbonyl carbon and the adjacent carbon

First phenyl hydrazone is formed and then reacts with an additional molecule of phenylhydrazine to form the osazones.

As is evident from the reaction, the first two carbons (C1 and C2) are involved in osazone formation

The sugars that differ in their configuration on these two carbons give the same type of osazones, since the difference is masked by binding with phenylhydrazine

Thus glucose, fructose and mannose give the same type (needle-shaped) osazones

Reducing disaccharides also give osazones

Maltose sunflower-shaped

And lactose powder puff shaped

Osazones have characteristic Melting points
Crystal structures
Precipitation time
and thus are used in identifying the sugars

9. Action of acids on carbohydrates

Polysaccharides are hydrolyzed into their constituent monosaccharides by boiling with **dilute** acids

With **concentrated** acids the monosaccharides are decomposed

Pentoses yield a cyclic aldehyde *furfural* with 12% HCL

Hexoses are decomposed by hot strong acids to give *hydroxy methyl furfural*

The furfural products thus formed can condense with certain organic (α-naphthol) to form compounds having characteristic colours
This forms basis of certain tests used for detection of sugars

The furfural products thus formed can condense with certain organic (α-naphthol) to form compounds having characteristic colours
This forms basis of certain tests used for detection of sugars

Molisch's and

Selivanoff's are examples of such tests

Disaccharides (such as maltose, lactose, and sucrose) consist of two monosaccharides joined covalently by an **O-glycosidic bond**.

Which is formed when a hydroxyl group of one sugar reacts with the anomeric carbon of the other

When the anomeric carbon is involved in **a glycosidic bond**, that sugar residue is not capable of reducing cupric ions and therefore becomes a non-reducing sugar

In describing disaccharides or polysaccharides, the end of a chain with a free anomeric carbon (one not involved in a glycosidic bond) is commonly called the reducing end

Maltose has two glucose units joined by an **α-1,4-glycosidic linkage**. It is a **reducing sugar** and also exhibits mutarotation

Maltose comes from the hydrolysis of starch (by **amylase**)
It is in turn hydrolysed to glucose by **maltase**, which is located on the intestinal brush border
From nutritional point of view, maltose is easily digestible
Various food preparations, such as baby and invalid foods, are produced by hydrolysis of grains and contain large amounts of maltose

Lesson 23 ""

Lactose, the disaccharide of milk consists of galactose joined to glucose by

"-6 9."glycosidic linkage

It is dextrorotatory

As one of the aldehyde group is free, it has reducing properties and exhibits mutarotation

In lactating mammary gland, the lactose is synthesized from glucose by the duct epithelium

Many organisms that are found in milk, e.g., *E.Coli* convert lactose of milk to lactic acid

(by **galactosidase**) thus causing souring of milk

Lactase, the specific enzyme which hydrolyses lactose is present in the intestinal brush border

Lactose intolerance refers to a condition of pain, nausea, and flatulence after the ingestion of foods containing lactose, most notably dairy products

Although it is often caused by

low levels of lactase

it also can because by intestinal injury

In a **lactase deficient** person, lactose accumulates in the lumen of the small intestine after ingestion of milk

The large osmotic effect of the unabsorbed lactose leads to an influx of fluid into the small intestine

Hence, the clinical symptoms of **lactose intolerance** are:

abdominal distention,

nausea,

cramping,

pain

a watery diarrhea

Lesson 24

Sucrose (Ordinary table sugar) is obtained commercially from cane or beet

The anomeric carbon atoms of a glucose unit and a fructose unit are joined in this disaccharide and the configuration of this glycosidic linkage is $\alpha(1\rightarrow2)\beta$

Consequently, sucrose lacks a free reducing group (an aldehyde or ketone end), in contrast with most other sugars

Sucrose is therefore a non-reducing sugar and does not exhibit mutarotation

Sucrose is dextrorotatory but its hydrolytic products are levorotatory as fructose has a levorotation greater than the dextrorotation of glucose

As the hydrolytic products invert the rotation, the resulting mixture is called as **invert sugar** and the process is called as **inversion**

The hydrolysis of sucrose to glucose and fructose is catalyzed by **sucrase** (also called **invertase**), which is also present in the intestinal brush border like lactase and maltase. Honey is largely invert sugar and the presence of fructose accounts for the greater sweetness of honey

Lesson 25

Oligosaccharides are short polymers of three to ten monosaccharides joined by glycosidic bonds.

Most are not digested by human enzymes

At one end of the chain, the reducing end, is a monosaccharide unit with its anomeric carbon not involved in a glycosidic bond

Biomedical Importance

Integral membrane proteins contain covalently attached carbohydrate units, oligosaccharides, on their extracellular face

Many secreted proteins such as antibodies and coagulation factors also contain oligosaccharide units

The oligosaccharides participate in molecular targeting and cell to cell recognition

They also mark the passage of time and determine when the proteins should be taken out of circulation

The human ABO blood groups illustrate the effects of glycosyl-transferases

Carbohydrates are attached to glycoproteins and glycolipids on the surfaces of red blood cells

For one type of blood group, one of the three different structures, termed A, B, and O, may be present

These structures have in common an oligosaccharide foundation called the O (or sometimes H) antigen

The A and B antigens differ from the O antigen by the addition of one extra monosaccharide, either *N*-acetylgalactosamine (for A) or galactose (for B) to the O antigen.

Lesson 26

Polysaccharides

Most carbohydrates found in nature occur as polysaccharides, polymers of medium to high molecular weight

Polysaccharides also called **glycans**, differ from each other

- in the identity of their recurring monosaccharide units
- in the length of their chains
- in the types of bonds linking the units
- in the degree of branching

Homopolysaccharides contain only a single monomeric species

Heteropolysaccharides contain two or more different kinds

Some **homopolysaccharides** serve as storage forms of monosaccharides that are used as fuels

Other homopolysaccharides (cellulose and chitin) serve as structural

elements in plants The most important storage polysaccharides are

starch in plant cells and glycogen in animal cells

Both polysaccharides occur intracellularly as large clusters or granules
Most plant cells have the ability to form starch and starch storage is especially abundant in tubers (underground stems), such as potatoes and in seeds

Heteropolysaccharides provide extracellular support for organisms of all kingdoms

For example, the rigid layer of the bacterial cell envelope (the peptidoglycan) is composed in part of a heteropolysaccharide built from two alternating monosaccharide units

Starch contains two types of glucose polymers:

Amylose
amylopectin

It is the most important dietary carbohydrate in cereals, potatoes, legumes, and other vegetables

Amylose (13–20%), consists of long, unbranched helical structure of D-glucose units linked by α -1,4 glycosidic bonds.

Lesson 27

Proteins mediate virtually every process that takes place in a cell, exhibiting an almost endless diversity of functions. To explore the molecular mechanism of a biological process, a biochemist almost inevitably studies one

or more protein

Proteins are the most abundant biological macromolecules occurring in all cells and all parts of cells

Proteins also occur in great variety; thousands of different kinds may be found in a single cell

Moreover, proteins exhibit enormous diversity of biological function and are the most important final products of the information pathways

Example signal transduction

Biochemical catalysts known as **enzymes** are proteins

Immunoglobulins that serve as the first line of defense against bacteria and viruses and other foreign agents are proteins

Several **hormones** are proteins

Structural proteins provide **mechanical support** e.g collagen

Contractile proteins help in the **movement** of muscle fiber and microvilli
Some proteins present in the cell membrane, cytoplasm and nucleus act as **receptors**

Certain other proteins in the cell membrane act as **channels and transporters**

The transport proteins carry out the function of transporting specific substances across the membrane or in the body fluids

Storage proteins bind with specific substances and store them

Under certain conditions, proteins can be catabolized to **provide energy**

Proteins help in the maintenance of water and electrolyte balance in the body by **exerting osmotic pressure**

Hence proteins are the most versatile macromolecules in living systems

and serve crucial functions in essentially all biological processes 

They function as catalysts, they transport and store other molecules such as oxygen,

They provide mechanical support and immune protection

They generate movement, and they control growth and differentiation

Lesson 28

Several key properties enable proteins to participate in such a wide range of functions

1. Proteins are linear polymers built of monomer units called amino acids

"The function of a protein is directly dependent on its three dimensional structure

"Remarkably, proteins spontaneously fold up into three-dimensional structures that are determined by the sequence of amino acids in the protein polymer

2. Proteins contain a wide range of functional groups

These functional groups include

alcohols,

thiols,

thioethers,

carboxylic acids,

carboxamides,

and a variety of basic groups

When combined in various sequences, this array of functional groups accounts for the broad spectrum of protein function

3. Proteins can interact with one another and with other biological macromolecules to form complex assemblies

The proteins within these assemblies can act synergistically to generate capabilities not afforded by the individual component proteins

These assemblies include macromolecular machines that carry out; the accurate replication of DNA,

the transmission of signals within cells, and many other essential processes

4. Some proteins are quite rigid, whereas others display limited flexibility

Rigid units can function as structural elements in the cytoskeleton (the internal scaffolding within cells) or in connective tissue
Parts of proteins with limited flexibility may act as hinges, springs, and levers

That are crucial to protein function, to the assembly of proteins with one another and with other molecules into complex units
And to the transmission of information within and between cells

Lesson 29

All proteins, whether from the most ancient lines of bacteria or from the most complex forms of life, are constructed from the same set of; 20 amino acids, "" Which are covalently linked in characteristic linear sequence.

Each of these amino acids has a side chain with distinctive chemical properties "" This group of 20 precursor molecules may be regarded as the alphabet in which the language of protein structure is written

Proteins are found in a wide range of sizes, from relatively small peptides with just a few amino acid residues to huge polymers with molecular weights in the millions
Cells can produce proteins with strikingly different properties and activities by joining the same 20 amino acids in many different combinations and sequence

From these building blocks different organisms can make such widely diverse products as

- enzymes,
- hormones,
- antibodies,
- transporters,
- muscle fibers,
- the lens protein of the eye,
- feathers,
- spider webs,
- rhinoceros horn,
- milk proteins,
- antibiotics,
- mushroom poisons etc

The light produced by fireflies is the result of a reaction involving the protein luciferin and ATP, catalyzed by the enzyme luciferase

RBC contain large amounts of the oxygen-transporting protein hemoglobin. (c) The protein keratin, formed by all vertebrates, is the chief structural component of hair, scales, horn, wool, nails, and feathers

Lesson 30

Proteins are polymers of amino acids, with each amino acid residue joined to its neighbor by a specific type of covalent bond termed the **peptide bond**

The term "residue" reflects the loss of the elements of water when one amino acid is joined to another

Twenty different amino acids are commonly found in proteins

All of the 20 common amino acids are **"j v rwx"j l m**

They have a carboxyl group and an amino group bonded to the same **nk°l yx" k' yw"- 'r o" "nk°l yx.**

They differ from each other in their side chains, or R groups, which vary in

structure,
size,
electric charge,
and which influence the solubility of the amino acids in water

Standard amino acids

The 20 amino acids that constitute the monomer units of proteins are **the standard amino acids** as the genetic code specifies only these 20 L - -amino acids

In addition to these 20 amino acids there are many less common ones i.e., **the non standard amino acids**

Lesson 31

Composition of Proteins

In addition to the 20 common amino acids, proteins may contain residues created by modification of common residues already incorporated in to a polypeptide

Composition of Proteins

Some proteins contain additional amino acids that arise by modification of an amino acid already present in a peptide i.e., after the protein has been synthesized

Examples include
conversion of peptidyl proline and lysine to
4-hydroxyproline and
5-hydroxylysine
the conversion of peptidyl glutamate to carboxyglutamate the methylation, formylation, acetylation, prenylation, and phosphorylation of certain aminoacyl residues

Among these uncommon amino acids

4-hydroxyproline, a derivative of proline

5-hydroxylysine, derived from lysine

The former is found in plant cell wall proteins, and both are found in collagen

6-N-Methyllysine is a constituent of myosin, a contractile protein

Another important uncommon amino acid is;

-carboxyglutamate, found in the blood clotting protein prothrombin and in certain other proteins that bind Ca^{2+} as part of their biological function

Some 300 additional amino acids have been found in cells but not as constituents of proteins

Examples include

Ornithine and citrulline (intermediates in urea cycle)

Taurine (found in bile acids)

-Aminolevulinic acid (intermediate in haem synthesis)

D- AMINO ACIDS

""D-amino acids are also non standard amino acids that occur **naturally** and include free D- serine, and D-aspartate in brain tissue

D-alanine & D-glutamate in the cell walls of gram positive bacteria

"" D-amino acids are also found in some **antibiotics**

Lesson 32

Selenocysteine, the 21st L- -Amino Acid?

Selenocysteine is an L- -amino acid found in a handful of proteins, including certain **peroxidases and reductases** where it participates in the catalysis of electron transfer reactions

Its synthesis is not a post translational modification, but a modification to a **serine** that occurs while the serine is bound to a unique RNA

The hydroxyl group of serine is replaced by a selenium atom

The selenocysteine is then inserted into a protein as it is being synthesized

Since selenocysteine is inserted into polypeptides during translation, it is commonly referred to as the "21st amino acid"

However, unlike the other 20 genetically encoded amino acids, selenocysteine is not specified by a simple three-letter codon

22nd Amino Acid

Pyrrolysine (abbreviated as Pyl or O) is a naturally occurring, genetically coded amino acid used by some methanogenic archaea.

serine

Lesson 33

Chiral centers and isomerism

"For all the common amino acids except glycine (the R group is another hydrogen atom), the central carbon atom is bonded to four different groups:

- a carboxyl group,
- an amino group,
- an R group,
- a hydrogen atom

The α -carbon atom is thus a chiral center and thus amino acids have two possible stereoisomers. All molecules with a chiral center are also optically active—that is, they rotate plane polarized light.

Special nomenclature has been developed to specify the absolute configuration of the four substituents of asymmetric carbon atom.

The absolute configurations of simple sugars and amino acids are specified by the D, L system, based on the absolute configuration of the three-carbon sugar glyceraldehyde.

Thus the amino group of L-alanine occupies the same position about the chiral carbon as does the hydroxyl group of L-glyceraldehyde.

Since they are nonsuperposable mirror images of each other, the two forms represent a class of stereoisomers called **enantiomers**.

Lesson 34

Two conventions are used to identify the carbons in an amino acid

The additional carbons in an R group are commonly designated **1', 1''** and so forth, proceeding out from the **1st**

For most other organic molecules, carbon atoms are simply numbered from one end, giving highest priority (C-1) to the carbon with the substituent containing the atom of **highest atomic number**

Within this latter convention, the carboxyl carbon of an amino acid would be **C-1** and the α carbon would be **C-2**

In some cases, such as amino acids with heterocyclic R groups (such as histidine), the

Greek lettering system is ambiguous and the numbering convention is therefore used

For branched amino acid side chains, equivalent carbons are given numbers after the Greek letters

So α carbons are $\alpha^1, \alpha^2, \alpha^3$ and β carbons are $\beta^1, \beta^2, \beta^3$

pros (in histidine nomenclature). The nitrogen atoms of the imidazole ring of
 r 3 'n3x0" k°o"noxy'on"l —z°y•"- xok° 1'kl l °o\3' on" . "kxn" "ow"- p k° 1'kl l °o\3' on"
 . "y•ryΔ" "r os"zy• s3yx"°ok's/o"y"r o"•no"m k3[r 3 "onywwoxnk'syx" k°y•o"
 from the fact that two different systems of numbering the atoms in the imidazole
 ring of histidine had both been used for a considerable time (biochemists
 generally numbering as 1 the nitrogen atom adjacent to the side chain, and
 organic chemists designating it as 3)

Lesson 35

Acid and base properties of amino acids

The form of an amino acid that has both a positive and a negative charge
 is called a **zwitterion**

Amino acids in solution at neutral pH exist predominantly as dipolar ions
 (**zwitterions**)

In this form, the amino group is protonated (NH_3^+)
 and the carboxyl group is deprotonated

(COO^-)

A zwitterion can act either as an acid (proton donor) or as a base (proton
 acceptor)

Substances having this dual nature are **amphoteric** and are often called
ampholytes

The ionization state of an amino acid varies with pH

At physiological pH, the carboxyl group is dissociated, forming the negatively charged carboxylate ion (-COO^-) and the amino group is protonated (NH_3^+)

A simple monoamino monocarboxylic - amino acid, such as alanine, is a diprotic acid when fully protonated. It has two groups, the -COOH group and the -NH_3^+ group, that can yield protons

Lesson 36

Each acid has a characteristic tendency to lose its proton in an aqueous solution

The stronger the acid, the greater its tendency to lose its proton

The tendency of any acid (HA) to lose a proton and form its conjugate base (A^-) is defined by the equilibrium constant (K_{eq}) for the reversible reaction



Equilibrium constants for ionization reactions are usually called ionization constants or acid dissociation constants, often designated pK_a

The stronger the tendency to dissociate a proton, **the stronger is the acid and the lower its pK_a**

The pK_a expresses, on a logarithmic scale, the relative strength of a weak acid or base:

'''

$$\text{pK}_a = \log$$

The stronger the acid, the lower its pK_a ;

'''

At the midpoint of the titration, the concentrations of the proton donor and proton acceptor are equal, and the pH is numerically equal to the pK_a .

The pK_a can be determined experimentally;

it is the pH at the midpoint of the titration curve for the acid or base

'''

At that point, $[\text{HA}]$ equals $[\text{A}^-]$, and

$$\text{pH} = \text{pK}_a + \log 1$$

$$= \text{pK}_a + 0 = \text{pK}_a$$

At that point, $[\text{HA}]$ equals $[\text{A}^-]$, and

$$\text{pH} = \text{pK}_a + \log 1$$

$$= \text{pK}_a + 0 = \text{pK}_a$$

Lesson 37

pK_a is the measure of the tendency of a group to give up (donate) a proton with the tendency decreasing ten-fold as the pK_a increases by one unit

Each titratable group has a pK_a that is numerically equal to the pH at which exactly one half of the protons have been removed from that group. The pK_a for the most acidic group ($COOH$) is pK_1 , whereas the pK_a for the next most acidic group (NH_3^+) is pK_2 .

In acid solution (e.g., pH 1), the amino group is protonated (NH_3^+) and the carboxyl group is not dissociated ($COOH$).

As the pH is raised, the carboxylic acid is the first group to give up a proton, as its pK_a is near 2.

The dipolar form persists until the pH approaches 9, when the protonated amino group loses a proton.

Lesson 38

Titration of an amino acid: Dissociation of the carboxyl group:

The titration curve of an amino acid can be analyzed.

At a low (acidic) pH, both of these groups are protonated. As the pH of the solution is raised, the $-COOH$ group of form I can dissociate by donating a proton to the medium.

The release of a proton results in the formation of the carboxylate group.

COO^-

This structure is shown as form II, which is the **dipolar form of the molecule**.

This form, also called a **zwitterion**, is the **isoelectric form** of alanine.

That is, it has an overall charge of zero.

The dissociation constant of the carboxyl group of an amino acid is called K_1 rather than K_2 because the molecule contains a second titratable group.

Dissociation of the amino group:

The second titratable group of alanine is the amino group.

This is a much weaker acid than the $-COOH$ group and, therefore, has a much smaller dissociation constant, K_2 .

Release of a proton from the protonated amino group of form II results in the fully deprotonated form of alanine, form III.

Lesson 39

A simple amino acid e.g. **glycine** is a diprotic acid when fully protonated. This means that it has two groups, the COOH and the NH_3^+ group that can yield protons.

Figure shows the titration curve of the diprotic form of glycine.

The two ionizable groups of glycine, the carboxyl group and the amino group, are titrated with a strong base.

The plot has two distinct stages, corresponding to deprotonation of two different groups on glycine.

At very low pH, the predominant ionic species of glycine is **the fully protonated form**,


$^+\text{H}_3\text{N}-\text{CH}_2-\text{COOH}$

At the midpoint of any titration, a point is reached where the **pH is equal to the pK_a** of the protonated group being titrated.

At the midpoint in the first stage of the titration, in which the -COOH group of glycine loses its proton, equimolar concentrations of the protonated ($^+\text{H}_3\text{N-CH}_2\text{-COOH}$) and the deprotonated ($^+\text{H}_3\text{N-CH}_2\text{-COO}^-$) species are present

As the titration proceeds, another important point is reached at pH 5.97. This is a point, at which removal of the first proton is essentially complete and removal of the second has just begun.


At this pH glycine is present largely as the dipolar ion (zwitterion)



The second stage of the titration corresponds to the removal of a proton from the NH_3^+ group of glycine.

The pH at the midpoint of this stage is 9.60, equal to the pK_a for the NH_3 group.

The titration is essentially complete at a pH of about 12, at which point the predominant form of glycine is



Lesson 40

From the titration curve of glycine we can derive several important pieces of information.

First, it gives a quantitative measure of the pK_a of each of the two ionizing groups:

2.34 for the -COOH group

9.60 for the NH_3^+ group

The second piece of information is that this amino acid has **two regions of buffering power**.

One of these is the relatively flat portion of the curve, extending for approximately 1 pH unit on either side of the first pK_a , of 2.6, indicating that glycine is a good buffer near this pH.

The other buffering zone is centered around pH 9.60.

Note that glycine is not a good buffer at the pH of intracellular fluid or blood, about 7.4.

Another important piece of information derived from the titration curve of an amino acid is the relationship between its net charge and the pH of the solution.

At pH 5.97, glycine is present predominantly as its dipolar form, fully ionized but with no net electric charge.

The characteristic pH at which the net electric charge is zero is called the **isoelectric point or isoelectric pH, designated pI**

For glycine, which has no ionizable group in its side chain; the pI is simply the arithmetic mean of the two pKa values

$$pI = \frac{1}{2} (pK_1 + pK_2)$$

$$= \frac{1}{2} (2.34 + 9.60) = 5.97$$

The farther the pH of a glycine solution is from its pI, the greater the net electric charge of the population of glycine molecules

At any pH below its pI, glycine has a **net positive charge** and will move toward the negative electrode (cathode)

At any pH above its pI, glycine will have a net negative charge and thus will move towards the positive electrode (anode)

Lesson 41

All amino acids with a

- -NH_2 -amino group,

- -COOH -carboxyl group, and

an R group that does not ionize

have titration curves resembling that of glycine

These amino acids have very similar, although not identical, pKa values:

pKa of the -COOH group in the range of 1.8 to 2.4

pKa of the -NH_3^+ group in the range of 8.8 to 11

The differences in these pKa values reflect the effects of the R groups

For example, the pKa of the -COOH group in glycine is 2.34, while in aspartic acid it is 1.88. The pKa of the -NH_3^+ group in glycine is 9.60, while in lysine it is 8.95.

Its pI can be calculated as follows:

$$pI = \frac{pK_1 + pK_2}{2}$$

$$pK_1 = 2.3 \text{ (for carboxyl group)}$$

$$pK_2 = 9.1 \text{ (for amino group)}$$

$$pI = \frac{2.3 + 9.1}{2} = 5.7$$

Amino acids with an ionizable R group have more complex titration curves, with three stages corresponding to the three possible ionization steps and thus they have **three pKa values**

The additional stage for the titration of the ionizable R group merges to some extent with the other two

In addition, each of the acidic and basic amino acids contains an ionizable group in its side chain

Thus, both free amino acids and some amino acids combined in peptide linkages can act as **buffers**

A buffer is a solution that resists change in pH following the addition of an acid or base

The isoelectric points reflect the nature of the ionizing R groups present

For example, glutamate has a pI of 3.22, considerably lower than that of glycine

This is due to the presence of two carboxyl groups, with an average of their pKa values

of

3.22

Histidine has got 3 dissociable hydrogens: one from carboxyl, $pK_1=1.8$, one from imidazole group, $pK_2=6.0$, one from amino group, $pK_3=9.2$

The pI of Histidine, with two groups that are positively charged when protonated, is 7.59 (the average of the pKa values of the amino and imidazole groups)

Only histidine has an R group ($pK_a = 6.0$) providing significant buffering power near the neutral pH usually found in the intracellular and extracellular fluids of most animals

42 Acidic and Basic amino acids

All amino acids with a

- -NH_2 -amino group,

- -COOH -carboxyl group, and

an R group that does not ionize

have titration curves resembling that of glycine

These amino acids have very similar, although not identical, pKa values:

pKa of the -COOH group in the range of 1.8 to 2.4

pKa of the -NH_3^+ group in the range of 8.8 to 11

The differences in these pKa values reflect the effects of the R groups

Alanine is an example of an amino acid with a non-ionizable R group. Its pKa values are 2.3 for the carboxyl group and 9.7 for the amino group.

Its pI can be calculated as follows:

$$\text{pI} = \text{pK}_1 + \text{pK}_2 \div 2$$

$$\text{pK}_1 = 2.3 \text{ (for carboxyl group)}$$

$$\text{pK}_2 = 9.7 \text{ (for amino group)}$$

$$\text{pI} = 2.3 + 9.7 \div 2 = 6.0$$

Amino acids with an ionizable R group have more complex titration curves, with three stages corresponding to the three possible ionization steps and thus they have three pKa values

The additional stage for the titration of the ionizable R group merges to some extent with the other two

In addition, each of the acidic and basic amino acids contains an ionizable group in its side chain

Thus, both free amino acids and some amino acids combined in peptide linkages can act as buffers

A buffer is a solution that resists change in pH following the addition of an acid or base

The isoelectric points reflect the nature of the ionizing R groups present

For example, glutamate has a pI of 3.22, considerably lower than that of glycine

This is due to the presence of two carboxyl groups, with an average of their pKa values of 3.22

Histidine has got 3 dissociable hydrogens

one from carboxyl, $pK_1=1.8$,

one from imidazole group, $pK_2=6.0$

one from amino group, $pK_3=9.2$

The pI of Histidine, with two groups that are positively charged when protonated, is 7.59 (the average of the pKa values of the amino and imidazole groups)

Only histidine has an R group ($pK_a = 6.0$) providing significant buffering power near the neutral pH usually found in the intracellular and extracellular fluids of most animals

43 Abbreviations and symbols of amino acids

Abbreviations and symbols for the commonly occurring amino acids

Each amino acid name has an associated three-letter abbreviation and a one-letter symbol

The one-letter codes are determined by the following rules:

1. Unique first letter:

If only one amino acid begins with a particular letter, then that letter is used as its symbol, for example,

I = isoleucine

2. Most commonly occurring amino acids have priority:

If more than one amino acid begins with a particular letter, the most common of these amino acids receives this letter as its symbol

For example, glycine is more common than glutamate,

So G = glycine

Similar sounding names:

Some one-letter symbols sound like the amino acid they represent

For example,

F = phenylalanine

Letter close to initial letter:

For the remaining amino acids, a one-letter symbol is assigned that is as close in the alphabet as possible to the initial of the amino acid

44 Classification of amino acids A. Aliphatic R groups

The topic can be simplified by grouping the amino acids into five main classes based on the properties of their R groups, in particular, their polarity, or tendency to interact with water at biological pH (near pH 7.0).

The polarity of the R groups varies widely, from nonpolar and hydrophobic (water-insoluble) to highly polar and hydrophilic (water-soluble)

A. Nonpolar, Aliphatic R Groups

These are hydrophobic

The amino acids in this group have non polar side chains that do not bind or give off protons

They also do not participate in hydrogen bonds

Zqn“”j °n"xnú‘ 'x°"úymłut n’j wrly°xv xµn"q“mfxyqxxkl 'hwµn°j l µxw " "

These side chains cluster together in the interior of the protein

This class includes

Glycine

Alanine

Proline

Valine

Leucine

Isoleucine

Methionine

Electrons are shared equally between the carbon and hydrogen atoms in these side chains, so that they can not hydrogen bond with water

Within proteins, these amino acid side chains will cluster together to form hydrophobic cores

45 Classification of amino acids A. Aliphatic R groups (Conti...)

Glycine has the simplest structure. Although it is formally nonpolar, its very small side chain makes no real contribution to hydrophobic interactions

It is the simplest amino acid, and it really does not fit well into any classification because its side chain is only a hydrogen atom

Because of this, it causes the least amount of steric hindrance in a protein

Therefore glycine is often found in bends or in the tightly packed chains of fibrous proteins.

Alanine and the branched chain amino acids (valine, leucine, and isoleucine) have bulky, nonpolar, *aliphatic(open chain hydrocarbon)* side chains

The side chains of alanine, valine, leucine, and isoleucine tend to cluster together within proteins, stabilizing protein structure by means of hydrophobic interactions.

Methionine, one of the two sulfur-containing amino acids, has a nonpolar thioether group in its side chain

The role of proline in amino acid structure differ from those of the nonpolar amino acids

The amino group of proline is part of the peptide backbone

It is an imino acid

This rigid ring causes a kink in the peptide backbone that prevents it from forming its usual configuration

Lesson 46

Classification of amino acids

B. Aromatic Amino Acids

The aromatic amino acids have been grouped together because they all contain ring structures with similar properties, **but their polarity differs**. The aromatic ring is a six-membered carbon hydrogen ring with three conjugated double bonds (the benzene ring or phenyl group). Benzene is an organic chemical compound with the chemical formula C_6H_6 . Its molecule is composed of 6 carbon atoms joined in a ring, with 1 hydrogen atom attached to each carbon atom. The substituents on this ring determine whether the amino acid side chain engages in polar or hydrophobic interactions. This group includes Phenylalanine, Tyrosine, Tryptophan, with their aromatic side chains, are relatively nonpolar (hydrophobic). All can participate in hydrophobic interactions.

In the amino acid **phenylalanine**, the ring contains no substituents, and the electrons are shared equally between the carbons in the ring, resulting in a very **nonpolar hydrophobic** structure in which the rings can stack on each other. In **tyrosine**, a hydroxyl group on the phenyl ring engages in hydrogen bonds, and the side chain is therefore more polar and **more hydrophilic**. The more complex ring structure in **tryptophan** is an indole ring with a nitrogen that can engage in hydrogen bonds. Tryptophan is therefore also more polar than phenylalanine. Tyrosine and tryptophan are significantly more polar than phenylalanine, because of the tyrosine hydroxyl group and the nitrogen of the **tryptophan** indole ring.

Lesson 47

C. Aliphatic, Polar, Uncharged R Groups

The R groups of these amino acids are more soluble in water, or more hydrophilic, than those of the nonpolar amino acids. Because they contain functional groups that form hydrogen bonds with water.

This class of amino acids includes **serine**, **threonine**, **cysteine**, " **asparagine**, and **glutamine**.

These Amino acids have zero net charge at neutral pH

Serine & threonine of this class contain a polar hydroxyl group. Similarly, side chains of **asparagine & glutamine** contain a carbonyl group & amide group, both of which can participate in hydrogen bonding.

These side chains are sites of attachment for other compounds.

The polar hydroxyl groups of **serine & threonine** serve as a site of attachment for phosphate groups.

The amide group of **asparagine** & the hydroxyl group of **serine & threonine** serve as a site of attachment for oligosaccharide chains in glycoproteins.

Amino acids with side chains that contain an amide group (asparagine and glutamine) or a hydroxyl group (serine and threonine) can be classified as **aliphatic, polar, uncharged amino acids**.

Asparagine and glutamine are amides of the amino acids aspartate and glutamate.

As a consequence of their hydrophilicity, these amino acids are frequently found on the surface of water-soluble globular proteins.

Cysteine, which is sometimes included in this class of amino acids, has been separated into the class of sulfur-containing amino acids.

Lesson 48

D. Positively Charged (Basic) R Groups

The side chains of these amino acids are proton acceptors.

The amino acids in which the R groups have significant **positive charge** at pH 7.0 are:

Lysine, which has a second α -amino group on its aliphatic chain.

Arginine, which has a positively charged guanidinium group.

Histidine, which has an aromatic imidazole group.

The side chains of the two basic amino acids, arginine and lysine, have pK_a values above 10, so that the positively charged form always predominates at physiologic pH.

The side chain of histidine ($pK_a \sim 6.0$) dissociates near physiologic pH, so only a portion of the histidine side chains carry a positive charge.

Histidine is therefore weakly basic & the *free* amino acid is mainly uncharged at physiologic pH.

The amino acid side chains might have very different pK_{as} than those of the free amino acids if they are involved in hydrogen or ionic bonds with other amino acid side chains

Therefore, when histidine is incorporated into a protein, its side chain can be either positively charged or neutral depending on the ionic environment provided by the polypeptide chains of the protein

Lesson 49

E. Negatively Charged (Acidic) R Groups

These amino acids are proton donors

At neutral pH, the side chains of these amino acids are fully ionized, containing a negatively charged carboxylate group

($-\text{COO}^-$)

Therefore these amino acids are negatively charged at physiologic pH

The two amino acids having R groups with a net negative charge at pH 7.0 are **aspartate and glutamate**, each of which has a second carboxyl group

Humans have no dietary requirement for protein, per se, but, the protein in food does provide essential amino acids. 

Nine of the twenty amino acids needed for the synthesis of body proteins are essential that is, they cannot be synthesized in humans at an adequate rate

Essential amino acids can not be made by the body. As a result, they must come from food. The nine **essential amino acids** are: histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine.

Of these nine, some amino acids are essential at all times, whereas some are required only during periods of rapid tissue growth characteristic of childhood or recovery from illness

Lesson 50

""Amino acids can be classified as **glucogenic or ketogenic** based on what intermediates are produced during their catabolism

A. Glucogenic amino acids

Amino acids whose catabolism yields **pyruvate** are termed **glucogenic or glycogenic**

These intermediates are substrates for **gluconeogenesis** and, therefore, can give rise to the net formation of glucose or glycogen in the liver and glycogen in the muscle

B. Ketogenic amino acids

Amino acids whose catabolism yields either **acetoacetate or its precursor, (acetyl CoA or acetoacetyl CoA)** are termed **ketogenic**

Acetoacetate is one of the ketone bodies which also include 3-hydroxybutyrate and acetone

Leucine and lysine are the only exclusively ketogenic amino acids found in proteins

Their carbon skeletons are not substrates for gluconeogenesis and, therefore, cannot give rise to the net formation of glucose or glycogen in the liver, or glycogen in the muscles

Lesson 51

""Different classifications of proteins are based on their:

1. Shape and Size
2. Biological actions/ Functions
3. Solubility and physical properties

4. **Quality Catalytic Proteins:**

These specialized proteins are called **enzymes** which catalyze the biochemical reactions

2. **Protective Proteins**

a) **Immunoglobins (Igs)**

These freely circulating proteins protect the body from invading microbes such as bacteria or viruses by inactivating or killing them through various mechanisms. **b) Fibrinogen**

This forms fibrin clot and stops bleeding from wounds

3. **Regulatory Proteins:**

Hormones control genetic expression, cellular signalling *and* biochemical reactions catalyzed by enzymes. Enzymes are either activated or inactivated through modification of their structure.

Examples of protein hormones are

growth hormone

Insulin

Glucagon Somatostatin

4. **Structural Proteins:**

These proteins form various body structures e.g.

Collagen

Elastin

Keratin

5. **Transport Proteins:**

These proteins transport various substances from one part of the body to the other e.g.

Hemoglobin transport O₂ from lungs to tissues and CO₂ from tissues to lungs Transferrin transports iron

6. **Contractile Proteins:**

These proteins are involved in muscle contraction and relaxation

Myosin of thick filaments

Actin of thin filaments of skeletal muscles

52 Functional classification of proteins

7. **Respiratory Proteins:**

"Heme containing proteins are involved in the function of respiration e.g.

Hemoglobin

Myoglobin

cytochromes

8. **Digestive Proteins:**

" These proteins are digestive enzymes which digest our food materials such as carbohydrates, proteins, lipids and include

Amylase

Pepsin

Lipases etc.

9. **Toxin Proteins:**

" These proteins are hydrolytic enzymes found in the venom of poisonous snakes, sting of bees and insects and hydrolyze the compounds forming the structure of the cell membrane

9. **Toxin Proteins:**

These proteins are hydrolytic enzymes found in the venom of poisonous snakes, sting of bees and insects and hydrolyze the compounds forming the structure of the cell membrane

The poisonous mushrooms also have such toxins

10. Storage Proteins:

These proteins store some specific elements or compounds with them

This is because of the presence of the many binding sites in them for the particular element e.g.

ferritin stores iron

ceruloplasmin stores copper

Lesson 53

PEPTIDE BOND

We now turn to polymers of amino acids, the peptides and proteins

Biologically occurring polypeptides range in size from small to very large, consisting of two or three to thousands of linked amino acid residues

Two amino acid molecules can be covalently joined through **an amide linkage**, termed **a peptide bond**

This linkage is formed by removal of the elements of water from the -NH_2 group of one amino acid and the -COOH group of another

Two configurations are possible for a planar peptide bond

In the trans configuration, the two α -carbon atoms are on opposite sides of the peptide bond

In the cis con-, these groups are on the same side of the peptide bond.

Almost all peptide bonds in proteins are trans

This preference for trans over cis can be explained by the fact that steric clashes between groups attached to the α -carbon atoms hinder formation of the cis form but do not occur in the trans configuration

Lesson 54

The Structure of Proteins

Four levels of protein structure are commonly defined

The Primary Structure

A description of all covalent bonds (mainly **peptide bonds** and **disulfide bonds**) linking amino acid residues in a polypeptide chain is its primary structure

The most important element of primary structure is the **sequence** of amino acid residues 

Secondary structure

Secondary structure refers to **particularly stable arrangements** of amino acid residues giving rise to recurring structural patterns

Tertiary structure

Tertiary structure describes all aspects of the **three-dimensional** folding of a polypeptide

Quaternary structure

When a protein has **two or more polypeptide** subunits, their arrangement in space is referred to as quaternary structure

Lesson 55 Primary structure

The primary structure of a protein refers to the **linear sequence** of amino acids in the polypeptide chain.

The primary structure is held together by covalent bonds such as peptide bonds, which are made during the process of protein biosynthesis or translation.

The two ends of the polypeptide chain are referred to as the **carboxyl terminus** (C-terminus) and the **amino terminus** (N-terminus) based on the nature of the free group on each extremity.

The primary structure of a protein determines **how it folds up** into its unique three-dimensional structure, and this in turn determines the function of the protein.

The function of a protein therefore depends on its amino acid sequence.

Proteins have unique amino acid sequences, that are specified by genes.

Each of the 20 amino acids incorporated in the proteins is encoded by one or more *specific* sequences of three nucleotides in DNA or RNA.

Understanding of the primary structure of proteins is important because many genetic diseases result in proteins with abnormal amino acid sequences which cause loss or impairment of normal function.

The defect can range from a single change in the amino acid sequence (as in **sickle cell anemia**) to deletion of a larger portion of the polypeptide chain (as in most cases of **Duchenne** muscular dystrophy).

In Sickle cell anemia a **Glutamate** residue is replaced by **Valine** at position 6 of beta chain of Hemoglobin, resulting in decreased functioning and increased fragility of Hb.

Duchenne Muscular dystrophy results from a large segment of gene deletion in X chromosome, resulting in small truncated **dystrophin** protein. Thus we know that if the primary structure is altered, the function of the protein may also be changed

Lesson 56

Secondary structure, the folding of short (3- to 30-residue), contiguous segments of polypeptide into **geometrically ordered units**

The term secondary structure refers to any chosen segment of a polypeptide chain and describes the local spatial arrangement of its main-chain atoms, without regard to its relationship to other segments

There are a few types of secondary structure that are particularly stable and occur widely in proteins

The most prominent are the **'α-helical' "**

conformations **β-sheets** **'β-barrel' "**

protein secondary structure

The simplest arrangement the polypeptide chain can assume, given its rigid peptide bonds (but free rotation around its other, single bonds), is a helical structure

where the carbonyl oxygen of one residue is hydrogen bonded to the amide hydrogen of a residue four positions further along the chain (the α-helix) participates in such hydrogen bonding

The structure is stabilized by a hydrogen bond

between the

hydrogen atom attached to the electronegative nitrogen atom of a peptide linkage and the electronegative carbonyl oxygen atom of the fourth amino acid on the amino-terminal side of that peptide bond

Lesson 57

Amino acid residues included in alpha helix have an **intrinsic propensity** to form an alpha helix, reflecting the properties of the R group.

Amino acids most commonly found in alpha helices are **non-polar** with aliphatic side chains

Alanine experimental model systems.

Amino acids most commonly found in alpha helices are **non-polar** with aliphatic side chains

Alanine experimental model systems.

The bulk and shape of **Ser**, **Thr**, and **Cys** helix if they are close together in the chain.

The bulk and shape of **Ser**, **Thr**, and **Cys** residues helix if they are close together in the chain.

Amino acids with charged R groups such as **Aspartate**, **Lysine** residues can also disrupt the helix.

Amino acids with charged R groups such as **Aspartate**, **Lysine** residues can also disrupt the helix.

In **proline**, the nitrogen atom is part of a rigid ring, and rotation about the N-C bond is not possible

[r ≠ 1k "z ° y v x o " o • n ≠ o " x ° y n ≠ m o • " k " n o • ' k l s s x q " u x u " x " k x " " r o v s - " "]

Glycine reason.

It has more conformational flexibility than the other amino acid residues

Polymers of glycine tend to take up cyson " ° ≠ m ≠ o • ~ ≠ s o " n s p o x " p y w " k x " " helix

Lesson 58 The Secondary Structure

A very diverse group of **α-helices**

The alpha-helical content of proteins ranges widely, from nearly none to almost 100%

For example, the **keratins** are a family of closely related, fibrous proteins that are α -helical.

They are a major component of tissues such as hair, nails, hoofs and skin.

Myoglobin is a globular, flexible molecule.

Nearly all trans membrane proteins contain alpha helices in their membrane spanning domains.

Lesson 59

Regular secondary structure that maximizes hydrogen bonding between the peptide backbones.

The backbone of the polypeptide chain is extended into a zigzag rather than helical structure

The zigzag polypeptide chains can be arranged side by side to form a structure resembling a series of pleats

Adjacent segments of polypeptide chain

The β -pleated sheet is described as parallel if the polypeptide strands run in the same direction (as defined by their amino and carboxy terminals) and anti-parallel if they run in opposite direction

Antiparallel strands are often the same polypeptide chain folded back on itself, with simple hairpin turns or long runs of polypeptide chain connecting the strands The amino acid side chains (R groups) of each polypeptide strand alternate between extending above and below the β -sheet. Most sheets are not perfectly flat but tend to have a twist.

β sheets are usually nearby on the polypeptide chain, but can also be quite distant from each other in the linear sequence of the peptide and may even be in different polypeptide chains.

Clusters of twisted strands of sheet form the core of many globular proteins.

Lesson 60

In contrast to regular repeating element in the twist of an alpha Helix or a pleat of Beta sheet, **bends, loops, and turns are** non-regular secondary structures that do not have a repeating element.

They often connect strands of antiparallel beta-sheets.

The structure is a 180 degree turn involving four amino acid residues

- Bends were given this name because they often connect successive

Nvns "kxn" Wyso "o•sn#o•" ypox"ym# "x" "≠rns

Proline because peptide bonds involving the imino nitrogen readily assume the cis configuration, a form that is particularly amenable to a tight turn

A **motif** also called a **super secondary structure** is simply a recognizable folding pattern involving two or more elements of secondary structure and the connection(s) between them.

A motif can be very simple, such as two elements of secondary structure folded against each other, and represent only a small part of a protein

A motif can also be a very elaborate structure involving scores of protein segments folded together, such as the 'kj'°°nu

Lesson 62

The tertiary structure

The tertiary structure of a protein is the folding pattern of the secondary structural elements into a three-dimensional conformation.

It indicates, in three-dimensional space, how secondary structural features helices, sheets, bends, turns, and loops assemble to form domains and how these domains relate spatially to one another.

Domains in the Tertiary Structure

The tertiary structure of large complex proteins is often described in terms of physically independent regions called structural domains.

A domain is a section of protein structure sufficient to perform a particular chemical or physical task such as binding of a substrate or other ligand.

Each domain is formed from a continuous sequence of amino acids in the polypeptide chain that are folded into a three-dimensional structure independently of the rest of the protein.

Two domains are connected through a simpler structure like a loop.

G-Actin. ATP binds in the center of the cleft. The two domains that form the cleft are further subdivided into subdomains 1–4. The overall structure is found in many ATP-binding proteins and is called the actin fold.

Lesson 63

The Tertiary Structure

Molecular forces that stabilise three dimensional tertiary structure include:

Electrostatic Interactions

Hydrophobic interactions

Disulphide bridges

Hydrogen bonding

The Tertiary Structure

The **positive** charges on the basic amino acids enables them to form ionic bonds (electrostatic bonds) with **negatively** charged groups, such as the side chains of acidic amino acids.

The Tertiary Structure

Electrostatic Interactions

The **positive** charges on the basic amino acids enables them to form ionic bonds (electrostatic bonds) with **negatively** charged groups, such as the side chains of acidic amino acids.

The Tertiary Structure

The Tertiary Structure

The positive charges on the basic amino acids enables them to form ionic bonds (electrostatic bonds) with negatively charged groups, such as the side chains of acidic amino acids.

The Tertiary Structure

The positive charges on the basic amino acids enables them to form ionic bonds (electrostatic bonds) with negatively charged groups, such as the side chains of acidic amino acids.

The Tertiary Structure

The positive charges on the basic amino acids enables them to form ionic bonds (electrostatic bonds) with negatively charged groups, such as the side chains of acidic amino acids.

The Tertiary Structure

Hydrophobic Interactions

Hydrophobic Interactions occur between non-polar uncharged side chains such as between aromatic groups of phenylalanine side chains.

The Tertiary Structure

Hydrophobic interactions occur between non-polar uncharged side chains such as between aromatic groups of phenylalanine side chains.

The Tertiary Structure **Disulfide Bridges**

The amino acid cysteine in a protein can form a covalent disulfide bond with another cysteine molecule through spontaneous (nonenzymatic) oxidation of their sulfhydryl groups.

The Tertiary Structure

The amino acid cysteine in a protein can form a covalent disulfide bond with another cysteine molecule through spontaneous (nonenzymatic) oxidation of their sulfhydryl groups.

The Tertiary Structure **Hydrogen bonds**

Hydrogen bonds in which a hydrogen atom is shared by a nitrogen (or oxygen) in the peptide backbone and an oxygen atom in an amino acid side chain.

The Tertiary Structure

Hydrogen bonds in which a hydrogen atom is shared by a nitrogen (or oxygen) in the peptide backbone and an oxygen atom in an amino acid side chain.

The Quaternary Structure

Some proteins contain two or more separate polypeptide chains, or subunits, which may be identical or different.

The arrangement of these protein subunits in three-dimensional complexes constitutes **quaternary structure**.

Many proteins have multiple polypeptide subunits (from two to hundreds)

The association of polypeptide chains can serve a variety of functions

A multisubunit protein is also referred to as a **multimer**, with the prefixes *homomultimer* or *heteromultimer* if the subunits are identical or different subunits, respectively.

Most multimers have identical subunits

or repeating groups of nonidentical subunits, usually in symmetric arrangements.

The repeating structural unit in such a multimeric protein is **protomer**.

Assembly into a multisubunit structure increases the stability of a protein.

X-ray diffraction analysis of deoxyhemoglobin shows how the four polypeptide subunits are packed together. (a) A ribbon representation. (b) A space-filling model. The alpha subunits are shown in gray and light blue; the beta subunits in pink and dark blue. Note that the heme groups (red) are  relatively far apart.

Actin filament is a homo polymer. G-actin, forms the basic unit for actin filaments. Actin filaments together with their associated motor proteins (e.g. myosin superfamily) form an elaborate network known as the actin cytoskeleton involved in functions including cell motility, muscle contraction, cell division, cytokinesis, vesicle and organelle movement and cell signaling.

Lesson 65

Perks of The Quaternary structure

The increase in size increases the number of possible interactions between amino acid residues and therefore makes it more difficult for a protein to unfold and refold.

A multisubunit structure has many advantages besides increased stability. It may enable the protein to exhibit cooperativity between subunits in binding ligands (illustrated later with hemoglobin)

Or it may form binding sites with a high affinity for large molecules (illustrated with antigen binding to the immunoglobulin molecule IgG).

An additional advantage of a multi-subunit structure is that the different subunits can have different activities and cooperate in a common function.

For example enzymes that have regulatory subunits or exist as multiprotein complex, such as Pyruvate Dehydrogenase.

E1 , pyruvate dehydrogenase;

E2, dihydrolipoyl transacetylase; and

E3,dihydrolipoyl dehydrogenase

(PDC) is a complex of three enzymes that convert pyruvate into acetyl-CoA by a process called pyruvate decarboxylation

Lesson 66 Protein Folding

Protein folding is the process by which a protein structure assumes its functional shape or conformation.

Protein Folding

It is the physical process by which a polypeptide folds into its characteristic and functional three-dimensional structure from random coil " Protein Folding

Protein Folding

The overall three-dimensional structure of a protein must meet certain requirements to enable the protein to function in the cell or extracellular medium of the body.

Protein Folding

The first requirement is the creation of a binding site that is specific for just one molecule, or a group of molecules with similar structural properties.

The specific binding sites of a protein usually define its functional role.

Protein Folding

Moreover, The three dimensional structure must have an external surface appropriate for its environment (e.g., plasma proteins contain polar amino acids on the surface to remain soluble in an aqueous environment.)

Protein Folding

Protein Folding

Proteins are dynamic molecules that can fold into their functionally competent conformation in milliseconds

Protein Folding

A protein is called a native protein if its amino acid composition and molecular conformation are unchanged from that found in natural states.

Protein Folding

Folding into the native state does not involve a haphazard search of all possible structures.

The native conformation of a Protein is dictated thermodynamically.

Protein Folding

The number of potential conformations of even a relatively small 15kDa polypeptide is unbelievably vast

"

Proteins are guided through this vast labyrinth of possibilities by thermodynamics

Protein Folding

Thermodynamic stability is not evenly distributed over the structure of a protein molecule.

It has regions of high and low stability; variations in the stability of

regions within a protein are often essential for protein to function. 

The regions of low stability allow a protein to alter its conformation between two or more states, such as an inactivated and activated enzyme.

Such as binding site for substrates in enzymes, ligands in receptors, for extracellular matrix in cytoskeletal proteins and for antigens in immunoglobins and so forth.

Lesson 67

Proteins are dynamic molecules that can fold into their functionally competent conformation in milliseconds

A protein is called a native protein if its amino acid composition and molecular conformation are unchanged from that found in natural states

And which is operative and functional

Folding into the native state does not involve a haphazard search of all possible structures.

Protein folding in cells takes place in an orderly and guided fashion. Protein folding generally occurs via a stepwise process.

In the first stage, as the newly synthesized polypeptide emerges from the ribosome, short segments fold into secondary structural units that provide local regions of organized structure

Then, each element of secondary or super-secondary structure facilitates proper folding by directing the folding process toward the native conformation and away from unproductive alternatives.

This step-wise folding of the proteins is dictated by thermodynamics resulting in a formation of native form, which is also the most energetically favoured form.

Computer simulated Protein folding showing *local assembly* followed by *global protein folding*.

Lesson 68

Protein Folding

The native conformation of a protein is dictated thermodynamically.

The biologically relevant or native conformation of a protein generally is that which is most energetically favored

The three dimensional structure of a native protein in its normal physiological state is the one in which the Gibbs free energy of the whole system is lowest.

The free energy change for a process at constant pressure is:

$$G = H - TS$$

G = Free Energy

H = Enthalpy

S = Entropy

Folding of a globular protein is a thermodynamically favored process, i.e. G must be negative.

$$G = H - TS$$

The folding process involves going from a multitude of random-coil conformations to a single folded structure.

The folding process involves a decrease in randomness and thus a decrease in entropy - S and an overall positive contribution to G. This decrease in entropy is

ΔS_{fold} = S_{folded} - S_{unfolded} < 0

Difference in energy (free energy) between folded (native) and unfolded (denatured) state is small, 5-15 kcal/mol.

Protein Folding

It is very difficult to determine how all factors blend together to give overall ΔG_{folding}

Use of averages contributions, but

Each protein is unique

Large stabilization factors,

Large destabilization factors,

But small difference between them

$$G = H - TS$$

ΔG_{fold} is necessary because too large a free energy change would mean a very stable protein, one that would never change.

However, structural flexibility is important to protein function, and proteins need to be degraded.

"

Enthalpy and entropy differences ΔH and ΔS are small positive numbers. ΔG is a small positive number.

Enthalpy is the total energy of the system. **Entropy** is a measure of the randomness or disorder in a system. **Entropy of a system always tends to increase to a maximum** value.

Lesson 69

Protein Folding

The native state has a smaller Gibbs free energy than the denatured state. The stability of the protein depends on the solvent-solvent, protein-solvent, and protein-protein interactions.

The free energy of a protein molecule is influenced by

- (1) the hydrophobicity,
- (2) hydrogen bonds,
- (3) electrostatic interactions, and
- (4) the conformational entropy due to the restricted motion

Entropy is a measure of the randomness or disorder in a system.

According to Second Law of Thermodynamics: **The entropy of a system will tend to increase to a maximum** value.

The folding process involves a decrease in randomness and thus a decrease in entropy $-\Delta S$ and an overall positive contribution to ΔG . This decrease in entropy is offset by the decrease in enthalpy $-\Delta H$ due to the formation of native contacts.

the hydrophobic effect and hydrogen bonds are the major stabilizing contributions which counteract entropy.

Starting from the unfolded protein, the polypeptide chain has to fold partially in order to bring together the residues that need to form the contacts stabilizing the native structure.

The constrained polypeptide chain has smaller entropy, which means higher Gibbs free energy.

As native contacts form, the enthalpy term decreases, the protein is stabilized.

Thermodynamic stability is not evenly distributed over the structure of a protein molecule.

It has regions of high and low stability; variations in the stability of regions within a protein are often essential for protein to function. 

The rate limiting step in the folding process is the formation of the transition state, i.e., the conformation that has the highest Gibbs free energy on the folding pathway

Entropy is maximum at unfolded state

The regions of low stability allow a protein to alter its conformation between two or more states, such as an inactivated and activated enzyme.

Lesson 70

Protein Folding

In the cell, not all proteins fold into their native conformation on their own.

As the protein folds and refolds while it is searching for its native low energy state, it passes through many high-energy conformations that slow the process

These high energy conformations are called kinetic barriers.

Cells employ auxiliary proteins to overcome these kinetic barriers.

These auxillary proteins are called chaperones.

Chaperone use energy provided by ATP hydrolysis to assist in the folding process

Chaperones participate in the folding of over half of mammalian proteins. They prevent immature folding of the nascent polypeptide and help in the final native protein formation.

The hsp70 family of chaperones binds short sequences of hydrophobic amino acids in newly synthesized polypeptides.

Protein Folding

Chaperones prevent aggregation, thus providing an opportunity for the formation of appropriate secondary structural elements.

Protein Folding

Protein Folding

They also unfold proteins prior to their insertion through the membrane of mitochondria and other organelles.

Lesson 71

Protein Folding

The hsp60 family of chaperones, sometimes called **chaperonins**, differ in sequence and structure from hsp70 and its homology

Hsp60 acts later in the folding process, often together with an hsp70 chaperone

Hsp60 chaperonins form a multi-subunit barrel-shaped structure

The unfolded protein fits into the barrel cavity that excludes water and serves as a template for the folding process.

Chaperone proteins can also "rescue" unfolded proteins that have become thermodynamically trapped in a misfolded dead end by unfolding hydrophobic regions and providing a second chance to fold productively

Finally, the folding pathways of some proteins require two enzymes that catalyze isomerization reactions.

Protein disulfide isomerase (PDI) is a widely distributed enzyme that catalyzes the interchange, or shuffling, of disulfide bonds until the bonds of the native conformation are formed.

The Hsp60 class of protein has a barrel shape into which the protein fits. It acts as a template, binding and rebinding portions of the unfolded protein until folding is completed. It hydrolyzes many ATP bonds to provide energy for the process.

Lesson 72

Protein Misfolding

Despite the many processes that assist in protein folding, misfolding does occur.

Misfolding of proteins may occur spontaneously

or caused by a mutation in a particular gene, producing an altered protein.

In addition some apparently normal proteins can, after abnormal proteolytic cleavage, take on a unique conformational state

[r k "vkn • "y"r o"p °wk 'yx"yp"yxq"p °sk °z °y °os "k • • owl v • "nyx • s 'sq"yp' " pleated sheets

Many conditions, including

Alzheimer's disease

Type 2 diabetes,

Huntington's disease and

Parkinson's disease, arise from a common misfolding mechanism.

In most cases, a soluble protein that is normally secreted from the cell is secreted in a misfolded state

and converted into an insoluble extracellular amyloid fiber.

The diseases are collectively referred to as amyloidoses.

Amyloidosis is a generic term that refers to the extracellular tissue deposition of fibrils composed of low molecular weight of a variety of proteins..

Amyloidoses may be

organ specific or generalized

(systemic).

Primary systemic amyloidosis is caused by deposition of fibrils consisting of misfolded immunoglobulin light chain

Examples of organ-specific amyloidoses include:

Gu'qnr v n° • 'm° nj • n'due to deposition of amyloid- β protein cleaved from amyloid precursor protein (APP).

Islet amyloid polypeptide (IAPP) is commonly seen in diabetes mellitus type 2 and is caused by deposition of amylin in pancreatic islets.

It has been suggested that the disease-causing mechanism in Huntington's disease (and the other polyglutamine disorders) is the ability of polyglutamine to undergo a conformational change that can lead to the formation of very stable anti-parallel betasheets; more specifically, amyloid structures

Photomicrograph of amyloid plaques in a section of temporal cortex from a patient with

Gu'qnr n° • 'h• nj • n

Alzheimer's is a type of dementia that causes problems with memory, thinking and behaviour. In August 1994, at the age of 83, Reagan was diagnosed with [Alzheimer's disease](#). Died in 2004.

73 Protein denaturation

Protein Denaturation

- A loss of three-dimensional structure sufficient to cause loss of function is called denaturation

(Electrostatic bonds, disulfide bonds, Specific hydrogen bonding, Hydrophobic bonds)

- The denatured state does not necessarily equate with complete unfolding of
µqn'y°xµrw] wml°j wmxv r'j µxw'xdl xwα°v j µxw'' '[wml°'v x• µl xwµµxw 1'
denatured proteins exist in a set of partially folded state

VU Bio Mates <https://www.facebook.com/groups/vubiomates/>

- When the urea and the reducing agent are removed, the randomly coiled, denatured ribonuclease spontaneously refolds into its correct tertiary structure, restoring its catalytic activity

VU Bio Mates

<https://www.facebook.com/groups/vubiomates/> denatur

renatur

Incubate 100-fold protein dilution of in protein guanidine

- Zqr 'y°xxo 'lqj plmwj µΔ°j µxw'xd' xv n'y°xµrw 'h°n.n° rkun3" "_ qrl q'pna "Δ•" that primary structure of a protein determines its three- dimensional conformation.

75 Reversible binding of a protein

Reversible binding of a protein to a ligand Oxygen-Binding

Proteins can bind to a variety of molecules, including oxygen, and this binding can be reversible.

binding of other molecules.

• The ability of a protein to bind to a ligand may be an essential part of its function. The nature of protein ligand interactions is critical to life, allowing an organism to respond rapidly and reversibly to changing environmental and metabolic circumstances.

• A ligand is a small molecule that is able to bind to proteins by weak interactions such as ionic bonds, hydrogen bonds, Van der Waals interactions, and hydrophobic effects. In some cases, a ligand also serves as a signal triggering molecule. A ligand can be a substrate inhibitor, activator or a neurotransmitter.

• A ligand binds at a site on the protein called the binding site, which is complementary to the ligand in size, shape, charge, hydrophobicity. This complementarity of protein ligand binding gives rise to the specificity of the binding. Hemoglobin's ability to bind and release oxygen is a classic example of reversible binding.

• The protein can discriminate among thousands of different molecules

in its environment and selectively bind only one or a few similar molecules. A protein may have separate binding sites for several different ligands. The binding of a ligand is often coupled to a conformational change in the protein that makes the binding site more complementary to the ligand, permitting tighter binding.

• The binding of a ligand to a protein can be reversible or irreversible. The binding of a ligand to a protein can be reversible or irreversible.

Specificity of ligands and binding sites. Ligand binding is often coupled to conformational changes in the protein. In multisubunit proteins, conformational changes in one subunit can affect the others (Cooperativity).

Examples of reversible binding include hemoglobin, antibodies, and muscle contraction.

76 Reversible binding of a protein (conti...)

Oxygen-Binding Proteins

- S “xpuκkw] wrlqnv xpuκkw” ‘Zqn n’l- x’v xul Δun ‘hΔ• p] j m’j w x• μh. n°“” aspect xdlqj μV x• μl nwμj d̄xdkxl qnv rl j d̄y°xl n• n• ?” ‘Zqn°n. n°• rkun” binding of j ‘l̄pj wrlx’j ‘y°xμrw” ‘U—“pnw’l ‘yx x°d̄”b xΔkun’w’j zΔnxΔ• ” solutions and cannot be carried to tissues in sufficient quantity if it is
- rv yu’‘h• • xu.nmlh/wkuxmlyj • v j 3” ‘J rΔ• ion of oxygen through tissues is also ineffective over distances greater than’j ‘t- ‘v ruv nμn°• ” ‘Zqn” evolution of larger, multicellular animals depended on the evolution of
- y°xμrw ‘lqj μl xΔunl̄j w yx°μj wrlb μx°n’x—“pnw” ‘S “xpuκkw’l ‘l xv yx• nml’ of a single polypeptide chain that has one O2 binding site.
- Hemoglobin is a tetramer in which each subunit has a strong sequence qxv xup“‘l’v “xpuκkw] wrl xwμj r• ‘j wU7’krmwμp’b n3” ‘S “xpuκkw] wrl’ hemoglobin are two oxygen-binding proteins with a very simila°” ‘y°rv j °”
- μΔl μΔ°n” ‘Nnv xpuκkw’l ‘j ‘l̄μj v n°l xv yx• nmlxd̄- x’hΔn°nwμtypes of
- ΔkΔwμp ‘-7 ‘j wrl7 ‘yxu’ynyμm’l qj r• 1°nΔ°nmlx’j • ‘l- x” ‘y°xμxv n°• ”
- Each subunit has a strong sequence homology to myoglobin.

- We will compare μqnA” ‘b μΔl μΔ°n’j wrl”
- functional relation of each protein in the subsequent discussion to see how both these proteins are suited to perform their required function in their relative sites in the body.

77 Reversible binding of a protein 1 (conti...)

Reversible binding of a protein

- The tertiary structure of oxygen binding guκkw ‘l xw r μ ‘xΔ” ‘Kpμμl qnul n• 3” ‘Zqn’qnul n• ‘l °nj m’j ” hydrophobic O2 binding pocket containing;
- tightly bound heme with an iron atom Ferrous (Fe 2+) in its centn°3” ‘63Muxkw373°x• μnμl ‘p°xΔy’l̄n’qnv n’l xv yx• nmlxd̄’ protoporphyrin IX rinp’l xwμj rrwμp’l’xw’j • ‘t°°xΔ• ” ‘Zqr ‘l’xw’kxΔwrl’ heme is an example of prosthetic group.

- A prosthetic group is a compound permanently associated with a protein and contributes to the protein's function.
- Prosthetic group is a tightly bound, specific non-polypeptide unit required for the biological function of some proteins. The prosthetic

group may be organic (such as a vitamin, sugar, or lipid) or inorganic (such as a metal ion), but is not composed of amino acids.

- Heme consists of a complex organic ring structure, protoporphyrin IX, to which is bound a single iron atom in its ferrous (Fe^{2+}) state

- Zqñ'f'xw'j µxv "qj • "b r-kxwm 1" "αΔ°"jx'wñfxpnw'j µxv • "lqj µj re part of µqn"qj µlyx°yq"°rw°rwp"b "• µnv "" "j wñlµ- x'yn°ynwml Δqj °"jx'lqn" porphyrin Negatively charged propionate (blue) groups on the porphyrin ring interact with arginine and histidine side chains from the hemoglobin, and the hydrophobic methyl(red) and vinyl(yellow) groups that extend out from the porphyrin ring interact with hydrophobic amino acid side chains from hemoglobin. All together, there are approximately 16 different interactions between myoglobin.

FIGURE 5 1 Heme. The heme group is present in myoglobin, hemoglobin, and many other proteins, designated heme proteins.

Heme

consists of a complex organic ring structure, protoporphyrin IX, with a bound iron atom in its ferrous (Fe^{2+}) state. µj µ3" Vx°yhyrins, of which protoporphyrin IX is only one example, consist of four pyrrole rings linked by methene bridges, with substitutions at one or more of the positions denoted X. (b, Two representations of heme (derived from PDB ID 1CCR). The iron atom of heme has six coordination bonds: four in the plane of, and bonded to, the flat porphyrin ring system, and (d) two perpendicular to it. Negatively charged propionate groups on the porphyrin ring interact with;

Arginine and

Histidine side chains from the hemoglobin. The hydrophobic; methyl and vinyl groups

- interact with hydrophobic amino acid side chains from hemoglobin.

- **78 Reversible binding of a protein 2 (conti...)**

- Iron in the Fe 2+ state binds oxygen reversibly

- The central iron atom of heme has six coordination bonds four in the plane of, and bonded to, the flat porphyrin ring system with

- The two axial bonds are to the proximal and distal ligands

- A coordinate bond (also called a dative covalent bond) is a covalent bond (a shared pair of electrons) in which both electrons come from the same atom.

Negatively charged propionate

(blue) groups on the porphyrin ring interact with arginine and

histidine side chains from the hemoglobin, and the hydrophobic

methyl (red) and vinyl (yellow) groups that extend out from the

porphyrin ring interact with hydrophobic amino acid side chains

from hemoglobin. All together, there are approximately 16 different interactions between myoglobin.

Heme:

protoporphyrin

- Therefore free heme molecules (heme not bound to protein) leave

Fe 2+ with

the proximal and distal ligands

• One of these two coordination bonds is occupied by a side chain
 Կապակցված է 'N-ի օրինակում՝ 'Z-ի օրինակում է 'կարգավորվող' դոմենում x-ի միջոցով
 oxygen (O) 2

• One of these two coordination bonds is occupied by a side chain
 Կապակցված է 'N-ի օրինակում՝ 'Z-ի օրինակում է 'կարգավորվող' դոմենում x-ի միջոցով
 oxygen (O) 2

Negatively charged propionate (blue) groups= arginine and histidine side

79 Reversible binding of a protein 3 (conti...)

Iron in the Fe^{2+} state binds oxygen reversibly

But in the Fe^{3+} (Ferric) state it does not bind oxygen

The coordinated nitrogen atoms (which have an electron-donating character) help in preventing the conversion of the heme iron to the ferric (Fe^{3+}) state

The conversion Fe^{2+} to Fe^{3+} is much more likely in *free* heme,

Sequestering each heme deep within the protein structure further helps to prevent this reaction.

Iron is a transition metal & because its orbitals are so close energy wise they tend to give up either 2 or 3 electrons at a time

Problem of Octet rule

Octet rule cannot be used for type II cations (most of the transition and inner transition elements).

The physiological oxidation state of an iron atom has a positive two charge (ferrous ion) instead of three charge (ferric ion)

And it is too large to fit into the plane of protoporphyrin

However, when iron oxidized from ferrous ion (Fe^{2+}) to ferric ion (Fe^{3+}), there is loss of one extra electron, forces between protons and electrons increases so that the electron cloud will penetrate more towards the nucleus

As a result, the ferric ion (Fe^{3+}) has a smaller size than ferrous ion (Fe^{2+}) and fits into the protoporphyrin plane

This deep impregnation of Fe^{3+} in the protoporphyrin results in inability of Fe^{3+} to bind with oxygen.

Some small molecules, such as carbon monoxide (CO) and nitric oxide (NO), coordinate to heme iron with greater affinity than does O_2

When a molecule of CO is bound to heme, O_2 is excluded, that is why CO is highly toxic to aerobic organisms

By surrounding and sequestering heme, oxygen-binding proteins regulate the access of CO to the heme iron.

80 Reversible binding of a protein 4 (conti..)

Helices with bends in between.

The helical segments are named A through H.(eight)

The bends are designated AB, CD, EF, and so forth.

An individual amino acid residue is designated either by its position in the amino acid sequence

or by its location in the sequence of a particular α -helical segment

For example, the His residue coordinated to the heme in myoglobin, His⁹³ (the 93rd residue from the amino-terminal end of the myoglobin polypeptide sequence), is also called His F8 (the 8th residue in a helix F)

Oxygen stored in red muscle myoglobin is released during O₂ deprivation (eg, severe exercise) for use in muscle mitochondria for aerobic synthesis of ATP.

The surface of myoglobin is polar, important for interacting with polar aqueous environment of cytosol.

The interior contains only nonpolar residues such as

Leu,

Val,

Phe, and

Met.

with two notable exceptions

The exceptions are His E7 and His F8, the seventh and eighth residues in helices E and F,

Which lie close to the heme iron, where they stabilize heme and help in O₂ binding. " Histidine is generally considered to be a *polar* amino acid

81 Reversible binding of a protein 5 (conti..)

The cervice (Pocket)created by nonpolar amino acids in the interior of myoglobin create a binding pocket for heme.

The proximal histidine F8, binds directly to the iron of heme

Whereas distal histidine E7, does not directly interact with the heme, but helps stabilize the binding of oxygen to the ferrous iron.

In myoglobin and hemoglobin, heme is covalently linked with histidine F8 (eighth residue of F helix). Because of covalent bond this histidine is closer to heme iron and named as proximal histidine (closer histidine),

While other key histidine which responsible for stabilization of oxygen in E7

(seventh residue of E helix) is far from heme iron so named as distal

The protein, or globin portion of myoglobin thus creates a special microenvironment for the heme that permits the reversible binding of one oxygen molecule (oxygenation)

The physiological importance of oxygen stabilization of the distal histidine (E7) lies in the fact that it prevents the binding of CO which is a low level product of cellular metabolism.

82 Reversible binding of a protein 6 (Conti....)

Structure and function of hemoglobin

Hemoglobin is found exclusively in red blood cells, where its main function is to transport oxygen from the lungs to the capillaries of the tissues

Hemoglobin A, the major hemoglobin in adults, is composed of four polypeptide chains

Two alpha - α chains

Two beta - β chains

HbF: In infants, made up of 2 alpha chains and 2 gamma chains. HbA₂: (alpha₂delta₂)

Two alpha chains and two beta chains are held together by hydrophobic interactions

The two polypeptide chains within each dimer are held tightly together, primarily by hydrophobic interactions

Inter- α chains are held together by hydrophobic interactions and hydrogen bonds. Inter- β chains are held together by hydrophobic interactions and hydrogen bonds. Inter- α and β chains are held together by ionic and hydrogen bonds.

Ionic and hydrogen bonds also occur between the members of the dimer

In contrast, the two dimers are able to move with respect to each other, being held together primarily by polar bonds

Each chain has a helical structure, with the heme group embedded in the

And a binding pocket similar to that described for myoglobin

Kj l q" ΔkΔwplbj • " μnp qn• "xδ" -helical structure,

And a binding pocket similar to that described for myoglobin

However, the tetrameric hemoglobin molecule is structurally and functionally more complex than myoglobin

For example, hemoglobin can transport CO₂ from the tissues to the lungs, and carry four molecules of O₂ from the lungs to the cells of the body

83 Reversible binding of a protein 7 (Conti....)

The proximal His (F8 histidine) of myoglobin and hemoglobin is sterically repelled by the heme porphyrin ring.

Thus, when the His binds to the Fe²⁺ in the middle of the ring, it pulls the Fe²⁺ above the plane of the ring.

The interior contains only nonpolar residues such as Leu, Val, Phe, Met. with exceptions of His (polar)

When oxygen binds on the other side of the ring, it pulls the Fe²⁺ back into the plane of the ring.

The pull of O₂ binding moves the proximal histidine toward the porphyrin ring, which moves the helix containing the proximal histidine.

This conformational change has no effect on the function of myoglobin.

However, in Hemoglobin this change is imperative for co-operative binding of oxygen to different subunits of hemoglobin.

Shifting of amino acids by the oxygenation of one of the heme groups in the protein, (due to pull of porphyrin through His), alters the structure of the interfaces between the four subunits. This results in the change of shape of the whole protein.

In the new shape, it is easier for the other three heme groups to become oxygenated.

Thus, the binding of one molecule of O₂ to hemoglobin enhances the ability of hemoglobin to bind more O₂ molecules.

This property of hemoglobin is known as "cooperative binding."

Cooperative binding of oxygen means that the binding of oxygen to one heme group enables an oxygen binding to the second heme group of the same hemoglobin molecule.

Conversely, the oxygen release from one heme group facilitates the release of oxygen from other heme groups in the same hemoglobin molecule.

84 Reversible binding of a protein 8 (Conti....)

In hemoglobin, the movement of one helix after oxygen binding leads to the movement of other helices in that subunit, including one in a corner of the subunit that is in contact with a different subunit through ionic interactions.

The loss of these ionic interactions, then induces conformational changes in all other subunits.

And all four subunits may change in a concerted manner from their original conformation to a new conformation structure bringing cooperativity in oxygen binding in hemoglobin.

As described previously, the binding of oxygen pulls the heme, which causes conformational change in the Polypeptide chain, resulting in the loss of some of the ionic interactions in the different subunits of Hb thus increasing the affinity of the Hb for the further O₂ addition.

Left: Hemoglobin Structure Showing Inter chain Contact Points Right : One of the contact points magnified, showing Lys, Asp and His.

Contact Points in Primary Structure: These amino acids which are far apart from each other in primary sequence become close in the final folding and give rise to characteristic interactions in oxygenated and unoxygenated states.

The conformational change of hemoglobin is usually described as changing from a T (tense) state with low affinity for O₂ to an R (relaxed) state with a high affinity for O₂.

In the T subunits, the binding sites are hindered "

And in the R state the binding sites are open.

Each successive addition of O₂ shifts the equilibrium further toward the R state, thus addition of oxygen in the lungs.

Similarly, removal of one oxygen facilitates the removal of subsequent oxygen molecule in the tissues.

85 Reversible binding of a protein 9 (Conti....)

Comparison of oxygen binding to myoglobin and hemoglobin

Myoglobin can bind only one molecule of oxygen.

Because it contains only one heme group

In contrast, hemoglobin can bind four oxygen molecules-one at each of its four heme groups

The degree of saturation (Y) of these oxygen-binding sites on all myoglobin or hemoglobin molecules can vary between zero (all sites are empty) and 100 percent

Oxygen dissociation curve

A plot of degree of saturation (Y) measured at different partial pressures of oxygen (pO_2) is called *the oxygen dissociation curve*

The curves for myoglobin and hemoglobin show important differences

This graph illustrates that myoglobin has a higher oxygen affinity than does hemoglobin

1 torr= 1mmHg

Furthermore, the hyperbolic versus sigmoid curve, in myoglobin and hemoglobin respectively, also shows that binding of oxygen is co-operative in hemoglobin as compared to myoglobin.

The *partial pressure of oxygen needed to achieve half-saturation of the binding sites is called (P_{50})*.

P_{50} is approximately 1 mm Hg for myoglobin and 26.6mm Hg for hemoglobin

The partial pressure of oxygen in the blood at which the hemoglobin is 50% saturated, is known as the P_{50} .

The P_{50} is a conventional measure of hemoglobin affinity for oxygen.

Values of P_{50} are negatively correlated with substrate affinity, with lower values of P_{50} corresponding to high affinity and *vice versa*.

This means that the higher the oxygen affinity (that is the more tightly oxygen binds), the lower the P_{50}

86 Reversible binding of a protein 10 (Conti....)

Myoglobin

The oxygen dissociation curve for myoglobin has a hyperbolic shape

"This reflects the fact that myoglobin reversibly binds a single molecule of oxygen

1 torr= 1mmHg Myoglobin

Myoglobin is designed to release and bind oxygen at very low partial pressure.

Oxygenated (MbO_2) and deoxygenated (Mb) myoglobin exist in a simple equilibrium:



The equilibrium is shifted to the right or to the left as oxygen is added to or removed from the system

Myoglobin is designed to bind oxygen released by hemoglobin at the low pO_2 found in muscle

Myoglobin, in turn, releases oxygen within the muscle cell in response to an oxygen demand during strenuous exercise.

When strenuous exercise lowers the pO_2 of muscle tissue to about 5 mm Hg, myoglobin releases O_2 for mitochondrial synthesis of ATP, permitting continued muscular activity

Myoglobin can load oxygen readily at the pO_2 of the lung capillary bed (100 mm Hg)

However, since myoglobin releases its bound O_2 only at the pO_2 values typically encountered in active muscle (20 mm Hg), it represents an ineffective vehicle for delivery of O_2 to tissues ($P_{O_2} = 40\text{mmHg}$)

87 Reversible binding of a protein 11 (Conti....)

Hemoglobin Dissociation Curve

Hemoglobin must bind oxygen efficiently in the lungs, where the pO_2 is about 100 mmHg,

And release oxygen in the tissues, where the pO_2 is about 35 to 40 mmHg

It is this sigmoidal loading and unloading of hemoglobin that makes it functionally useful as an oxygen transporter in blood.

Sigmoidal or sigmoid, literally means S-shaped

The sigmoidal binding curve of Hemoglobin for oxygen is possible due to multisubunit structure of hemoglobin. i.e

$\mu\text{-x}^j \text{ yqj } ^{-} .^j \text{ wr}^n$

$\mu\text{-x}^k \text{ nj } ^{-} .^l \text{ qj } \text{rw}^n$

As described earlier binding of one O_2 molecule induces conformational change in hemoglobin structure resulting in transition of hemoglobin from the low-affinity T (taut) state to the high-affinity R (relaxed) state

These changes significantly increase the affinity of the remaining unoxygenated hemes for O₂, as subsequent binding require the disruption of fewer ionic interactions.

The transition from T to R does not take place after a fixed number of oxygen molecules have been bound

But becomes more probable as each successive oxygen binds

Every oxygen molecule that binds to Hb increases its affinity for the binding of further oxygen molecules.

This effect is referred to as heme-heme interaction

88 Reversible binding of a protein 12 (Conti....)

Hemoglobin Dissociation Curve

Although it is more difficult for the first oxygen molecule to bind to hemoglobin.

The subsequent binding of oxygen occurs with high affinity, as shown by the steep upward curve in the region near 20 to 30 mm Hg

Cooperative binding of oxygen by the four subunits of hemoglobin means that the binding of an oxygen molecule at one heme group increases the oxygen affinity of the remaining heme groups in the same hemoglobin molecule

The net effect is that the affinity of hemoglobin for the last oxygen bound is approximately 300 times greater than its affinity for the first oxygen bound

In Summary,

Hb must bind O_2 efficiently in the lungs ($pO_2 = 100$ mm Hg), and release O_2 in the tissues ($pO_2 = 40$ mm Hg).

Hb solves the problem by undergoing a transition from a low-affinity state (the T state) to a high-affinity state (the R state) as more O_2 molecules are bound. "

Hb has a hybrid S-shaped, or sigmoid, binding curve for O_2

89 Reversible binding of a protein 13 (Conti....)

Allosteric effects

Apart from Oxygen, ability of hemoglobin to reversibly bind oxygen is affected by

- the pH of the environment

- the pCO_2 ; and

- the availability of 2,3-bisphospho-phoglycerate

These are collectively called Allosteric (other site) effectors

because their interaction at one site on the hemoglobin molecule affects the binding of oxygen to heme groups at other locations on the molecule

In Greek, Allo means other and Steric means Site.

The binding of oxygen to myoglobin is not influenced by the allosteric effectors of hemoglobin

In addition to transporting O_2 from the lungs to the peripheral tissues,

hemoglobin transports CO_2 , the by product of respiration,

And protons from peripheral tissues to the lungs

Transport of Carbon Dioxide

Hemoglobin carries CO_2 as carbamates formed with the amino terminal nitrogens of the polypeptide chains.

Carbamates change the charge on amino terminals from positive to negative, favoring the stabilization of the taut form of deoxyhemoglobin

Hence, favoring the stabilization of taut form of deoxyhemoglobin

Hemoglobin carbamates account for about 15% of the CO₂ in venous blood

The remainder of the CO₂ is transported as dissolved

HCO₃⁻ (80 to 85%) and
dissolved CO₂ (<5%)

Modes of CO₂ transport

1: Dissolved CO₂; 2: Dissolved bicarbonate 3: carbamate form

90 Reversible binding of a protein 14 (Conti....)

Transport of Carbon Dioxide (cont.)

Much of the remaining CO_2 is carried as bicarbonate.

(80 to 85%)

Bicarbonate is formed in erythrocytes by the hydration of CO_2 to carbonic acid (H_2CO_3)

This process is catalyzed by carbonic anhydrase

At the pH of venous blood, H_2CO_3 dissociates into;

bicarbonate and

a proton

Deoxyhemoglobin binds one proton for every two O_2 molecules released, contributing significantly to the buffering capacity of blood

Oxygen and H^+ are not bound at the same sites in hemoglobin.

However binding of proton, helps stabilize deoxyhemoglobin in the T state.

The somewhat lower pH of peripheral tissues, aided by carbamation, stabilizes the T state and thus enhances the delivery of O_2

This reciprocal coupling of proton and O_2 binding is termed the Bohr effect

91 Reversible binding of a protein 15 (Conti....)

This reciprocal coupling of proton and O_2 binding is termed the Bohr effect

Chloride Shift: When bicarbonate is formed, chloride ions move into the RBCs, to maintain

electroneutrality This is referred to as Chloride shift.

In the lungs, the process reverses

As O_2 binds to deoxyhemoglobin, protons are released and combine with bicarbonate to form carbonic acid

Dehydration of H_2CO_3 catalyzed by carbonic anhydrase forms CO_2 , which is exhaled

Binding of oxygen thus drives the exhalation of CO₂

92 Reversible binding of a protein 16 (Conti....)

Bohr effect

Hemoglobin transports about 40% of the total H⁺

And 15% to 20% of the CO₂ formed in the tissues to the lungs and the kidneys.

Bohr effect

Hemoglobin transports about 40% of the total H⁺

And 15% to 20% of the CO₂ formed in the tissues to the lungs and the kidneys.

Bohr effect

The binding of H⁺ and CO₂ is inversely related to the binding of oxygen.

Bohr effect

At the relatively low pH and high CO₂ concentration of peripheral tissues, the affinity of hemoglobin for oxygen decreases as H⁺ and CO₂ are bound, and O₂ is released to the tissues.

Bohr effect

Conversely, in the capillaries of the lung, as CO_2 is exhaled

Bohr effect

And the blood pH consequently rises, the affinity of hemoglobin for oxygen increases and the protein binds more O_2 for transport to the peripheral tissues.

Bohr effect

This effect of pH and CO_2 concentration on the binding and release of oxygen by hemoglobin is called the Bohr effect, after Christian Bohr, the Danish Physiologist who discovered it in 1904

93 Reversible binding of a protein 17 (Conti....)

OxyHemoglobin Dissociation Curve

Bohr effect decreased affinity of hemoglobin for oxygen caused by an increase of carbon dioxide pH etc.

OxyHemoglobin Dissociation Curve

Therefore oxyhemoglobin dissociation curve is displaced to the right because of higher partial pressure of carbon dioxide and lower pH.

OxyHemoglobin Dissociation Curve

This curve describes the relationship between available oxygen and amount of oxygen carried by hemoglobin.

The horizontal axis is PaO_2 , or the amount of oxygen available.

The vertical axis is SaO_2 , or the amount of hemoglobin saturated with oxygen.

Once the PaO_2 reaches 60 mm Hg the curve is almost flat, indicating there is little change in saturation above this point.

So, PaO_2 of 60 or more is usually considered adequate.

But, at less than 60 mm Hg the curve is very steep, and small changes in the PaO_2 greatly reduce the SaO_2 .

The term "affinity" is used to describe oxygen's attraction to hemoglobin binding sites. Affinity changes with:

variation in

pH,

temperature,

CO₂ and,

2,3,-BPG

These are the metabolic by-products which competes with O₂ for binding sites.

Traditionally the curve starts with:

pH at 7.4,

temperature at 37 Centigrade, and

PaO₂ at 40.

Changes from these values are called "shifts

94 Reversible binding of a protein 18 (Conti....)

Bohr effect

Hemoglobin transports about 40% of the total H⁺

And 15% to 20% of the CO₂ formed in the tissues to the lungs and the kidneys.

Bohr effect

The remainder of the H⁺ is transported by bicarbonate ions.

And the remainder of the CO_2 is transported as dissolved HCO_3^- and CO_2 .

Bohr effect

The binding of H^+ and CO_2 is inversely related to the binding of oxygen.

Bohr effect

At the relatively low pH and high CO_2 concentration of peripheral tissues, the affinity of hemoglobin for oxygen decreases as H^+ and CO_2 are bound, and O_2 is released to the tissues.

Bohr effect

Conversely, in the capillaries of the lung, as CO_2 is exhaled

Bohr effect

And the blood pH consequently rises, the affinity of hemoglobin for oxygen increases and the protein binds more O_2 for transport to the peripheral tissues.

Bohr effect

This effect of pH and CO_2 concentration on the binding and release of oxygen by hemoglobin is called the Bohr effect, after Christian Bohr, the Danish Physiologist who discovered it in 1904

95 Reversible binding of a protein 19 (Conti....)

Proton Binding (Cont.)

In the tissues, as the concentration of H^+ rises, (i.e. decrease in pH), protonation of His HC3 occurs.

This protonation promotes release of oxygen by favoring a transition to the T state, subsequently resulting in oxygen unloading.

In the lungs,

the pH of blood is higher because CO_2 is being exhaled,

And hemoglobin is being oxygenated due to higher partial pressure of oxygen in alveoli.

The increase in pH,

exhalation of CO_2

and oxygenation of Hb;

all result in; deprotonation of His HC3 of Hb.

This disrupts the ionic interactions that stabilize the taut state

And a new stabilized state of relaxed Hb is formed, resulting in oxygen loading.

Some ion pairs that stabilize the T state of deoxyhemoglobin: Interactions between the ion pairs His HC3 and Asp FG.1 of the Beta subunit and between Lys C5 of the alpha Subunit and His HC3 of the Beta subunit are shown with dashed line.

Binding of O₂, protons and CO₂ to the Hb is a dynamic process, in which relative concentrations of each of the compounds in the different compartments of the body dictate the binding of each other.

Thus we have seen that the four polypeptide chains of hemoglobin communicate with each other about not only O₂ binding to their heme groups

But also H⁺ binding to specific amino acid residues

96 Reversible binding of a protein 20 (Conti....)

Proton Binding (cont.)

An increase in protons and/or a higher pCO₂ shifts the equilibrium to the right

Therefore favoring deoxyhemoglobin T form.

Whereas a decrease in pCO₂ and/or a decrease in protons shifts the equilibrium to the left

Conversely, an increase in pO₂ promotes proton release

Decreased pH and increased CO₂ shift the curve to right, i.e. . *increased* O₂ unloading : Increased pH and decreased CO₂ shift the curve to left i.e. *decreased* O₂ unloading.

Decreased pH and increased CO₂ shift the curve to right, i.e. *increased* O₂ unloading : Increased pH and decreased CO₂ shift the curve to left i.e. *decreased* O₂ unloading.

The differential pH gradient (lungs having a higher pH, tissues a lower pH) favors the unloading of oxygen in the peripheral tissues,

And the loading of oxygen in the lung

In Summary,

Binding of oxygen to facilitate other molecule of oxygen is called positive cooperativity.

Whereas, binding of carbon dioxide and proton to release oxygen is called negative allostery, in which;

increased CO₂ and H⁺ ion favor oxygen unloading.

Thus we see that the four polypeptide chains of hemoglobin communicate with each other about not only O₂ binding to their heme groups

But also H⁺ binding to specific amino acid residues and CO₂ binding to amino terminal.

It is the capacity to communicate ligand binding information from one polypeptide subunit to the others that makes the hemoglobin molecule so beautifully adapted to integrate the transport of O₂, CO₂ and H⁺ by RBCs.

And there is still more to the story.

Many more interactions of different amino acids, are being identified as new techniques are employed to study these effects, which are out of the scope of this course.

97 Reversible binding of a protein 21 (Conti....)

2,3 bisphosphoglycerate

The interaction of 2,3 bisphosphoglycerate (BPG) with hemoglobin further refines the function of hemoglobin

and provides example of heterotropic allosteric modulation.

2, 3 BPG is an alternate product of glycolysis and its concentration increases in RBCs in states of low oxygen delivery to the tissues.

It is a negatively charged molecule which increases the stability of taut form of hemoglobin.

2,3-Bisphosphoglycerate is known to greatly reduce the affinity of hemoglobin for oxygen.

"
"

There is an inverse relationship between the binding of O₂ and the binding of BPG.

BPG binds at a site distant from the oxygen-binding site

And therefore regulates the O₂-binding affinity of hemoglobin in relation to the pO₂ in the lungs.

BPG is important in the physiological adaptation to the lower pO₂ at high altitudes

At sea level, the binding of oxygen to hemoglobin is regulated such that the amount of oxygen delivered to the tissues is nearly 40% of the maximum that could be carried by the blood

At high altitude of hills and mountains, where the pO₂ is considerably lower (due to low atmospheric pressure).

The delivery of oxygen to the tissues is now reduced.

However, after just a few hours at the higher altitude, the BPG concentration in the blood begins to rise, leading to a decrease in the affinity of hemoglobin for oxygen.

This adjustment in the BPG level has only a small effect on the binding of oxygen in the lungs

But a considerable effect on the release of oxygen in the tissues shown by increased oxygen unloading at the peripheral tissues.

Increase in BPG concentration in the RBC shifts the dissociation curve to the right, i.e. increased O₂ unloading.

BPG concentration also increases in those conditions in which there is decreased O₂ delivery to tissues, such as in anemia and respiratory diseases.

As a result, the delivery of oxygen to the tissues is restored to nearly 40% of the oxygen that can be transported by the blood.

98 Reversible binding of a protein 22 (Conti....)

2,3 bisphosphoglycerate

(Count.)

The site of BPG binding to Hb is the cavity between the beta subunits.

This cavity is lined with positively charged amino acid residues that interact with the negatively charged groups of BPG.

"
"

BPG forms ionic interactions with the terminal amino groups of both chains via Val NA1 and with Lys EF6 and His H21.

NA is a non-helical portion between the start of amino terminal and start of A helix of beta chain.

Val NA1 is the first amino acid in this non-helical portion.

NA is a non-helical portion between the start of amino terminal and start of A helix of beta chain.

Val NA1 is the first amino acid in this non-helical portion.

BPG therefore stabilizes deoxygenated (T state) hemoglobin by forming additional ionic interactions that must be broken prior to conversion to the R state.

Unlike oxygen, only one molecule of BPG is bound to each hemoglobin tetramer

BPG lowers hemoglobin's affinity for oxygen by stabilizing the T state

The transition to the R state narrows the binding pocket.

Therefore BPG, precluding BPG binding.

Xn• mΔn'N76'xδlqn" " ΔkΔwplxδNkL is Ser rather than His which cannot form ionic interactions with BPG

Therefore, BPG binds more weakly to HbF than to HbA.

The lower stabilization afforded to the T state by BPG accounts for HbF having a higher affinity for O₂ than HbA

99 Reversible binding of a protein 23 (Conti....

Sickle Cell Anemia

Sickle Cell Anemia is a molecular disease of hemoglobin.

It demonstrates the importance of amino acid sequence in hierarchical structure of globular proteins, and thus their biological functions

Sickle cell anemia is a genetic disorder caused by a single nucleotide alteration (a point mutation) w'lqn" -globin gene.

This results in the production of altered hemoglobin.

The point mutation in the DNA sequence result in the replacement of; Glutamate residue with Valine residue

Glyx• rpxw'; 'xδlqn" 'l qj w'xδqnv xpuxkwβ'''

Hnl j Δ• n'qΔv j w 'pnwn°j w'qj .n'l- x'l xym• 'xδ' -globin gene.

An individual may have two copies of mutant allele (i.e. homozygous) or one copy of each of normal and mutant allele

(i.e. heterozygous)

Sickle-cell anemia occurs in individuals who inherit the allele for sickle-cell hemoglobin from both parents.

Tqn°n• Δpwp'qnv xpuxkw' 2 s₂, is referred to as HbS

The RBCs of these individuals are fewer and also abnormal.

In addition to an unusually large number of immature cells, the blood contains many long, thin, sickle shaped RBCs.

When HbS is deoxygenated, it becomes insoluble and forms polymers that aggregate into tubular fibers.

The insoluble fibers of deoxygenated hemoglobin S cause the deformed, sickle shape of the RBCs.

The proportion of sickled cells increases greatly as blood is deoxygenated.

Normal hemoglobin (hemoglobin A) remains soluble on deoxygenation.

100 Reversible binding of a protein 24

Sickle Cell Anemia

(Count.)

The altered properties of hemoglobin S result from the replacement of Val
 ᎠᏍᏉᏅ ᎡᏂᏗᏛ 'ᎢᏰᏱᏁ ᎡᏂᏗᏛ' ᎡᏂᏗᏛ ᎡᏂᏗᏛ; 'ᎡᏂᏗᏛ' ᎡᏂᏗᏛ ᎡᏂᏗᏛ

The R group of valine is neutral, whereas glutamate has a negative charge at pH 7.4.

Hemoglobin S therefore has two fewer negative charges than hemoglobin A.

Replacement of the Glu residue by Val creates a "sticky" hydrophobic
 I xwuj l μyxrwuj μ'yx• rpxw', 'xdlqn" "l qj rwf– qrl q'x' 'xwujqn'xΔμ°b Δ°g l n'xδ'
 the molecule

These sticky spots cause deoxyhemoglobin S molecules to associate abnormally with each other,

Therefore forming the long, fibrous aggregates characteristic of this disorder.

The polymerization of Hb S inside the cytosol of RBCs make RBCs sickle shaped and fragile..

Normal Hb A does not have this sticky hydrophobic pocket and therefore is soluble in the cytosol.

The lifetime of an RBC homozygous for HbS is approximately 20 days, compared with 120 days for normal red blood cells.

Sickled cells are very fragile and rupture easily; this results in anemia.

Moreover, these cells have less ability to squeeze through small blood vessels resulting in the occlusion of micro blood vessels in the body, i.e. capillaries.

Sickle-cell anemia, occurs in individuals homozygous for the HbS allele,

Individuals who receive the HbS allele from only one parent (i.e. heterozygous) experience a milder condition called sickle-cell trait.

Only about 1% of their RBCs become sickled on deoxygenation.

101 Lipid Chemistry

The lipids are a heterogeneous group of compounds, including fats, oils, steroids, waxes,

their derivatives

and precursors,

that are related more by their physical than by their chemical properties.

They have the common property of being

(1) Insoluble in water

(2) Soluble in nonpolar solvents such as ether and chloroform.

Lipids are characterized by their hydrophobicity and their immiscibility in water

The biological functions of the lipids are as diverse as their chemistry.

Fats and oils are the principal stored form of energy in many organisms.

Phospholipids and sterols are major structural elements of biological membranes.

Other lipids, present in relatively small quantities, play crucial roles as; enzyme cofactors, electron carriers, hydrophobic anchors for proteins, emulsifying agents in the digestive tract, hormones, and cellular messengers

Adipose tissue Fat also serves as a thermal insulator in the subcutaneous tissues and around certain organs.

Nonpolar lipids act as electrical insulators, allowing rapid propagation of depolarization waves along myelinated nerves

Myelination with FAT greatly reduce the loss of charges from nerve fibre, resulting in a High speed transmission of nerve impulse.

The fat-soluble vitamins serve as co enzymes for many important reactions in the body.

Combinations of lipid and protein (lipoproteins) serve as the means of transporting lipids in the blood

Importantly, lipids provide the hydrophobic barrier that permits partitioning the aqueous contents of cells and subcellular structures as; phospholipids and sterols are the major structural elements of biological membranes.

102 Lipid Chemistry Classification of lipids

CLASSIFICATIONS OF LIPIDS

As mentioned previously, lipids are characterized by their hydrophobicity.

The precursors and derivatives of such compounds are included in classification of lipids.

Lipids are classified in following three major groups:

1. Simple lipids
2. Complex Lipids and
3. Precursor and Derived Lipids

1. Simple lipids:

Esters of fatty acids with various alcohols

These contain:

- a. Fats (and Oils) and
- b. Waxes.
- c. a. Fats: Esters of fatty acids with glycerol (Oils are fats in the liquid state)
- d. b. Waxes: Esters of fatty acids with higher molecular weight monohydric alcohols.
- e. (having one OH group)

2. Complex lipids: Esters of fatty acids containing groups in addition to an alcohol and a fatty acid.

" These include:

- a) Phospholipids
- b) Glycolipids
- c) Other Complex lipids
- d) a. Phospholipids: Lipids containing, in addition to fatty acids and an alcohol, a phosphoric acid residue
- e) Phospholipids frequently have nitrogen containing bases and other substituent:
- f) Glycerophospholipids the alcohol is glycerol.
- g) Sphingophospholipids the alcohol is sphingosine.
- b. Glycolipids (glycosphingolipids):
Lipids containing a;
fatty acid,
sphingosine, and
carbohydrate
- C. Other complex lipids: Lipids such as
sulfolipids

aminolipids

lipoproteins may also be placed in this category

3. Precursor and derived lipids:

" These include

fatty acids

glycerol

steroids

other alcohols

fatty aldehydes

ketone bodies


lipid-soluble vitamins

103 Lipid Chemistry Fatty Acids

Fatty Acids

The fats and oils used almost universally as stored forms of energy in living organisms are derivatives of fatty acids

Fatty acids are aliphatic carboxylic acids with hydrocarbon chains.



Basic structure of Fatty Acid: Long chain of hydrocarbon terminating in the carboxylic group.

Hexa: 6; Deca: 10

Hexadecanoic acid: 16 carbon Fatty acid.

Fatty acids occur in the body mainly as esters in natural fats and oils,

But are found in the unesterified form as free fatty acids, a transport form in the plasma.

Fatty acids that occur in natural fats usually contain an even number of carbon atoms ranging from 4 to 36 carbons (C₄ to C₃₆)

Fatty acids are named after the corresponding hydrocarbons.

A fatty acid consists of

a hydrophobic hydrocarbon chain with

a terminal carboxyl group that has a pKa of about 4.8

At physiologic pH, the terminal carboxyl group (-COOH) ionizes, becoming COO^-

This anionic group has an affinity for water, giving the fatty acid its amphipathic nature

(having both a hydrophilic and a hydrophobic region)

However, for long-chain fatty acids (LCFA), the hydrophobic portion is predominant

These molecules are highly water-insoluble, and must be transported in the circulation in association with protein.

More than ninety percent of the fatty acids found in plasma are in the form of;

fatty acid esters (primarily triacylglycerol, cholesteryl esters, and phospholipids) contained in circulating lipoprotein particles

Triacylglycerol: An example of Fatty acid containing compound, R denotes the hydrocarbon chain.

Unesterified fatty acids are transported in the circulation in association with albumin.

Low levels of free fatty acids occur in all tissues,

But substantial amounts sometimes can be found in the plasma, particularly during fasting.

104 Lipid Chemistry Classification Of Fatty Acids

Classification Of Fatty Acids:

Fatty Acids are classified on the basis of :

Hydrocarbon Chain Length

Degree of Saturation

Dietary Requirement

Classification on the basis of dietary requirement include essential and non-essential fatty acids.

Classification On The Basis Of Hydrocarbon Chain Length

These include

Short Chain Fatty Acids (2-4 C)

Medium Chain Fatty Acids (6-12 C)

Long Chain Fatty

Acids (14-18 C)


Very Long Chain Fatty Acids (18 and more C)

Saturation of fatty acids

Fatty acid chains may contain

no double bonds, that is be saturated, or

contain one or more double bonds that is, be mono- or polyunsaturated



Oleic acid 18-carbon

When double bonds are present, they are nearly always in the cis rather than in the trans configuration.

If the fatty acid has two or more double bonds they are always spaced at three carbon intervals.

Oleic acid 18-carbon, monounsaturated

Linoleic acid 18-carbon, polyunsaturated

Classification of Fatty Acid on Dietary Basis

Fatty acids that are required by the human body but

cannot be synthesized in the body, and

therefore must be obtained from food, are called essential fatty acids. 

In contrast to amino acids, there are only two essential Fatty Acids:

Linoleic Acid

Alpha Linolenic Acid

Humans lack the ability to introduce double bonds in fatty acids beyond carbons 9 and 10 as counted from the carboxylic acid side.

Linoleic Acid has two double bonds at C9 and C12.

γ-linolenic acid has three double bonds at C 9, 12 and 15.

Therefore, both of these fatty acids can not be synthesized in the body and must be obtained in the diet.

The rest of the Fatty Acids fall into the category of non-essential fatty Acids.

105 Lipid Chemistry Nomenclature of Fatty Acids

Nomenclature

The most frequently used systematic nomenclature names the fatty acid after the hydrocarbon with the same number and arrangement of carbon atoms, with

-oic being substituted for the final e

Thus, saturated acids end in -anoic. Octane becomes Octanoic Acid.

Unsaturated acids with double bonds end in -enoic, e.g. octadecenoic acid (C18) also called oleic acid.

A simplified nomenclature (more aptly called as symbolic nomenclature) for unbranched fatty acids specifies

the chain length and

number of double bonds,

separated by a colon.

The carbon atoms are numbered, beginning with the carboxyl carbon as C 1

" The number before the colon indicates the total number of carbons in the chain,

Those after the colon indicate the numbers and positions of double bonds.

The 16-carbon saturated palmitic acid is abbreviated

16:0,

The 18-carbon oleic acid, with one double bond, is

18:1

The carbon to which the carboxyl group is attached, carbon 2 is also called α carbon

l j °kxw'8'r "μgn" -carbon, and

l j °kxw'9'ኮ '፲፱n" -carbon

The positions of any double bonds are specified relative to the carboxyl carbon by superscript numbers follx- rwp" "rmuj ."

A 20-carbon fatty acid with two double bonds between C-9 and C-10 (C-1 being the carboxyl carbon) and another between C-12 and C-13 is designated " **757**" ^{9,12}

106 Lipid Chemistry Nomenclature of Fatty Acids (Conti....)

The family of polyunsaturated fatty acids (PUFAs) with a double bond between the third and fourth carbon from the methyl end of the chain are of special importance in human nutrition

Arachidowl 'j l m759 5,8,11,14 r °nœ°nm|œ'j • 'j w' -6 fatty acid because the
l œ • œ hœΔkun'kœwmlœ'œœn' 'hwmlknprœ'b œ—j œkœns from that end.

Zgn'h•nwuj uq̄ μ'j | ml'wuxuml 'j | ml'6=?" ^{9, 12} r" -6 fatty acid.

Q1 xwuj • uj' -uwxunwl 'j l ml'6=3 9, 12, 15 r 'j w' -3 fatty acid

107 Lipid Chemistry Physical and Chemical Properties of Fatty Acids
Physical and Chemical Properties of Fatty Acids

These are dictated by

length of hydrocarbon chain and

degree of saturation and unsaturation of Fatty Acids.

The melting points T_m of even-numbered carbon fatty acids increase with increasing chain length and

decrease with increasing unsaturation.

A triacylglycerol containing three saturated fatty acids of 12 carbons or more is solid at body temperature, whereas

if the fatty acid residues are 18:2, i.e. unsaturated it is liquid to below 0°C

Carbon Atoms: Double Bonds	Common Name	Melting Point (°C)
Saturated Fatty Acids		
12:0	Lauric acid	44
14:0	Myristic acid	58
16:0	Palmitic acid	63
18:0	Stearic acid	70
20:0	Arachidic acid	77
Unsaturated Fatty Acids		
16:1	Palmitoleic acid ¹	
18:1	Oleic acid	16
18:2	Linoleic acid	-5
18:3	Linolenic acid	-11
20:4	Arachidonic acid	-49

At room temperature (25°C), the saturated fatty acids from 12:0 to 24:0 have a waxy consistency, whereas

unsaturated fatty acids of these lengths are oily liquids.

Increasing chain length increases the melting temp. Stearic acid, Oleic acid, Linoleic acid and Linolenic acid all have 18 C, but with increasing no of Double bonds, which decreases the T_m

This difference in melting points is due to different degrees of packing of the fatty acid molecules due to;

different proportions of saturated and unsaturated Fatty Acids

Saturated fatty acids are more tightly packed as compared to "

Unsaturated Fatty Acids; which have a less compact packing.

The more compact packing;

the more attractive molecular forces between Fatty Acid molecules and vice versa.

Hence, saturated Fatty Acids have high T_m and

unsaturated Fatty Acids have low T_m.

Therefore saturated and Unsaturated Fatty Acids help to Shape Foods

108 Lipid Chemistry Physical and Chemical Properties of Fatty Acids (Conti....)

Physical and Chemical Properties of Fatty Acids (Contd.)

The melting points (Contd.)

The loose packing of unsaturated fatty acids is due to; cis isomerism of unsaturated fatty acids.

Unsaturated fatty acids show

Geometric isomerism, i.e.

They are either

cis or

trans

When double bonds are present, they are nearly always in the cis rather than in the trans configuration.

A cis configuration means that the two hydrogen atoms adjacent to the double bond stick out on the same side of the chain.

Unsaturated fatty acids can be cis with bulky groups on same side of C=C.


Unsaturated fatty acids can be cis with bulky groups on same side of C=C.



"



trans have bulky groups on opposite sides of C=C.



"



trans



Cis configuration leads to kink, which causes loose packing, have low T_m.

cis configuration, the molecules bent 120 degrees at the double bond. Trans 180 degree

In unsaturated fatty acids, a cis double bond forces a kink in the hydrocarbon chain

Fatty acids with such kinks

cannot pack together as tightly as fully saturated fatty acids,

resulting in weak interaction with each other.

In practice, natural acylglycerols contain a mixture of fatty acids tailored to suit their functional roles

The membrane lipids, which must be fluid at all environmental temperatures, are more unsaturated than storage lipids.

Saturated; All the carbons on the fatty acid are bound to hydrogen. Solid at room temp.

Higher melting point

Trans fatty acids are present in certain foods, arising as a by-product of the saturation of fatty acids during q“nfxpnrwj μxwfx°" qj °mwrwp1 'xδWj μΔ°j ḁXrṁ "hẉlqn" manufacture of margarine.

An additional small contribution comes from the ingestion of ruminant fat that contains *trans* fatty acids arising from the action of microorganisms in the rumen.

109 Lipid Chemistry Physical and Chemical Properties of Fatty Acids 1(Conti....) Physical and Chemical Properties of Fatty Acids (Contd.)

Water Solubility

The presence of nonpolar hydrocarbon chain results in the poor water solubility of fatty acids.

The larger the fatty acyl chain

and the fewer the double bonds,

the lower is the solubility in water.

The carboxylic acid group is polar and

ionized at neutral pH and

accounts for the slight solubility of

short-chain fatty acids in water

Melting Points and Solubility in Water of Fatty Acids

Formation of Esters

In combination with alcohol Fatty Acids form Esters.

For example, fatty acids react with glycerol to form mono, di and tri glycerides.

110 Lipid Chemistry Physical and Chemical Properties of Fatty Acids 2 (Conti....)

Physical and Chemical Properties of Fatty Acids (Contd.)

Formation of Salts

Fatty Acids forms salts with alkali metals and alkaline earth metals.

Salts of Na, K, Ca and Mg are called soaps.

A soap micelle: nonpolar (hydrophobic) hydrocarbon chains cluster in the inside and polar (hydrophilic) carboxylate groups lie on the surface.

The Na and K salts are water soluble,

Whereas, Ca and Mg salts are insoluble.

K salts are too soluble and are used in liquid soap preparations.

Formation of Detergents

Reduction of the carboxyl group of fatty acids produce alkyl alcohols

which can be sulfated (OSO_3H) or sulfonated (SO_3H) to produce detergents.

Lauric acid or dodecanoic acid, the saturated fatty acid with a 12-carbon atom chain, thus falling into the medium chain fatty acids

111 Lipid Chemistry Special Reactions of Unsaturated Fatty Acids

Special Reactions of Unsaturated Fatty Acids

Hydrogenation

It is the addition of Hydrogen at the double bonds of unsaturated fatty acids.

Hydrogenation converts

Unsaturated fatty acids to " Saturated fatty acids.

Margarines are vegetable oils

treated with partial hydrogenation to

form semi- $\bullet$ xumll xw r μwl “"kΔμh° "

Hydrogenated fats are used by many commercial food producers to

Provide rich texture

Increase shelf live

Increase melting point

Resistance to oxidation and flavour deterioration

Partial hydrogenation has an undesirable, effect:

some cis double bonds are converted
to trans double bonds.

There is a strong evidence that dietary intake of trans fatty acids
(often referred to as "trans fats")

leads to a higher incidence of cardio-vascular disease,
therefore these fats should be avoided in the diet.

Halogenation

Similar to hydrogen,

Halogens such as chlorine, bromine and iodine can also be added to double bonds in

unsaturated fatty acids.

Degree of halogenation is a

good index of

degree of unsaturation of Fatty Acids

The number of grams of iodine which will be absorbed by 100 grams of a fat is termed its iodine number.

112 Polyunsaturated Fatty Acids of Biological Importance

Lipid Chemistry

Polyunsaturated Fatty Acids Of Biological Importance "Polyunsaturated

fatty acids contain more than one double bond

These include:

Monounsaturated fatty acids (9, 12)

-Monounsaturated fatty acids (9, 12, 15)

Essential fatty acids (5, 8, 11, 14)

Of these

Linoleic acid and

-Linolenic acid are essential fatty acids. 

Linolenic acid is a polyunsaturated fatty acid with three double bonds. It is an essential fatty acid, meaning it cannot be synthesized by the body and must be obtained from the diet.

Arachidonic acid is the precursor of paracrine hormones called Eicosanoids.

Eicosanoids are chemical messengers which act over a short distance and include:

Prostaglandins

Leukotrienes and

Thromboxanes

These eicosanoids are semi cyclic structures.

They exert complex control over many bodily systems; in

inflammation

immunity, and as

messengers in the central nervous system.

FINAL TERM 113 TO 225

BIO504T - Biochemistry-I

114 Polyunsaturated Fatty Acids of Biological Importance

Polyunsaturated fatty acids contain more than one double bond

These include:

Stearic acid (18:0)

- Stearic acid (18:0)

Arachidonic acid (20:4)

Of these

α-Linolenic acid are essential fatty acids. 

Whereas, arachidonic acid is synthesized from α-Linolenic acid.

Arachidonic acid becomes essential if linoleic acid is deficient in the diet

"

'''

Arachidonic acid is the precursor of paracrine hormones called **Eicosanoids**.

Eicosanoids are chemical messengers which act over a short distance and include:

Prostaglandins

Thromboxanes

These eicosanoids are semi cyclic structures.

They exert complex control over many bodily systems; in inflammation immunity, and as messengers in the central nervous system.

115 Polyunsaturated Fatty Acids Of Biological Importance Contd.

Polyunsaturated Fatty Acids Of Biological Importance (Cont.)

Eicosanoids are derived from either omega-3 or omega-6 fatty acids.

Omega-6 eicosanoids are pro-inflammatory.

Figure 2. Classes of Essential Fatty Acids





Figure 2. Classes of Essential Fatty Acids

Omega-3 fatty acids

- Alpha-Linolenic acid (ALA) Green leafy vegetables Flax and chia seeds
- Stearidonic acid (SDA) Canola, walnut, and Soybean oils
- Eicosatetraenoic acid (ETA)
- Eicosapentaenoic acid (EPA) Oily fish Algae oil Krill oil
- Docosapentaenoic acid



" **Imbalance of omega-6 and omega-3 PUFAs** in the diet is associated with:
an increased risk of cardiovascular disease.

Long chain omega-3 fatty acids such as alpha-linolenic acid and their derivatives such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) have anti-inflammatory effects. In addition, docosahexaenoic acid (DHA) is selectively incorporated into retinal cell membranes and postsynaptic neuronal cell membranes suggesting it plays important roles in vision and nervous system function. Current evidence suggests that diet rich in omega 3 fatty acids are beneficial particularly for cardiovascular disease and also to some extent for Cancer and Rheumatoid Arthritis. This disease preventing role is due to the anti-inflammatory action of these omega-3 derivatives. Omega-3 PUFAs mediate some of these effects by inhibiting omega-6 PUFA (arachidonic acid)-induced pro-inflammatory prostaglandin E (PGE) formation.

116 Unsaturated Fatty Acids of Physiologic and

Nutritional Significance

Linseed .Alsi borage oil: Gao Zaban primrose: Gul e Fanjani (used in skin care products)

117 Lipid Chemistry-Other Common Fatty Acids

Other Common Fatty Acids

Other Fatty acids commonly found in body which are worth knowing are:

Palmitic Acid (16:0)

Stearic Acid (18:0) and Oleic acid (18:1)

Palmitic Acid(16:0) Palmitic acid, or hexadecanoic acid, is the most common saturated Fatty Acid found in animals, plants and microorganisms " Palmitic acid mainly occurs as its ester in triglycerides (fats), especially palm oil.

It is also found in high amounts in

Butter,

Cheese,

milk and

meat

Excess carbohydrates in the body are converted to palmitic acid.

Palmitic acid is the first fatty acid produced during fatty acid synthesis and the precursor to longer fatty acids

As a consequence, palmitic acid is a major body component of fats found in the animals.

Stearic Acid (18:0)

As its ester, stearic acid is one of the most common saturated fatty acids found in nature following palmitic acid.

Fats and oils rich in stearic acid are more abundant in animal fat (up to 30%) than in vegetable fat (typically <5%)

Oleic Acid(18:1)

Possibly the most common fatty acid in adipose tissue " It is particularly high in olive oil

Oleic acid in olive oil accounts for the blood pressure lowering effect of olive oil. " In addition, because of cis configuration and the kink produced due to the cis configuration.

It is also highly abundant in membrane lipids, providing fluidity to membrane.

Saturated Fatty Acids

Saturated Fatty Acids

Spermaceti by the sperm whale. Palm saeed. evergreen tree (*Laurus nobilis*) of the Mediterranean region valued for its aromatic ovate leaves. Nutmeg Jorenk, Jefal

118 Lipid Chemistry-Glycerol and Sphingosine

Glycerol and Sphingosine are the two types of alcohols most commonly found in lipids.

"N^oy"

It is a simple poly hydroxy alcohol (also called polyol or sugar alcohol)

and part of a class of lipids: glycolipids

It contains 3 carbons and 3 hydroxyl (OH) groups.

Glycerol is synthesized from

Dihydroxyacetone Phosphate (an intermediate of the glycolytic pathway)

Dihydroxyacetone phosphate is acted upon by two enzymes to form glycerol, namely Glycerol Phosphate Dehydrogenase and " Glycerol Kinase respectively.

Glycerol is a precursor for synthesis of triacylglycerols and of phospholipids in the liver and adipose tissue.

When the body uses stored fat as a source of energy, glycerol and fatty acids are released into the bloodstream.

And enter the glycolysis pathway directly

119 Lipid Chemistry: Sphingosine

Sphingosine is an amino alcohol,

which is a component of the class of lipids known as sphingolipids " Sphingosine is synthesized in the body in the form of ceramide,

to which different moieties are added to form sphingolipids.

Serine and palmitoyl CoA condense to form a product (ketosphinganine) that is reduced.

A very long-chain fatty acid forms an amide with the amino group.

a double bond is generated, and ceramide is formed.

There is no direct route of synthesis from sphinganine (dihydrosphingosine) to sphingosine;
it has to be acylated
first to dihydroceramide, which is then dehydrogenated to ceramide.
Sphingosine is formed via degradation of sphingolipid in the lysosome.
Therefore ceramide is the structural parent of all sphingolipids.

In short,
Serine+ Palmitate
Sphingosine
Sphingosine +
FA = ceramide

120 Lipid Chemistry-Properties of Glycerol

Popularly known as glycerin,

Glycerol is widely used in pharmaceutical and cosmetic preparations.

It has the following properties:

Colorless Viscous oily liquid with sweet taste.

Acrolein Test

On heating with sulfuric acid or KHSO_4 (dehydration)

it gives acrolein that has a bad odor.

used for detection of free glycerol or any compound containing glycerol.

In contrast to glycerol

Sphingosine does not show positive acrolein test.

Therefore glycerolipids and shingolipids can be differentiated on the basis of acrolein test.

Glycerol combines with three molecules of nitric acid to form Glycerol trinitrate that is used as explosive and vasodilator

On esterification with fatty acids it gives:

monoacylglycerol: one fatty acid + glycerol.

diacylglycerol: two fatty acids + glycerol.

triacyl-glycerol: three fatty acids + glycerol.

Gluconeogenic substrate Glycerol can form dihydroxyacetone phosphate which can convert into glucose or glyceraldehyde phosphate to enter into gluconeogenic pathway.

121 Lipid Chemistry-Simple lipids

Esters of fatty acids with various alcohols

These contain:

Fats (and Oils) and Waxes.

Fats: Esters of fatty acids with glycerol (Oils are fats in the liquid state)

Waxes: Esters of fatty acids with higher molecular weight monohydric alcohols.
(having one OH group)

Triacylglycerols (TAGs)

The simplest lipids constructed from fatty acids are the triacylglycerols,

Also referred to as;

triglycerides,

fats, or neutral fats or storage lipids.

Triacylglycerols are composed of three fatty acids in ester linkage with a single glycerol

Structure of triacylglycerols

The three fatty acids esterified to a glycerol molecule are usually not of the same type

The fatty acid on carbon 1 is typically saturated,

and the fatty acid on carbon 3 is typically unsaturated.

Simple triglycerides:

Fatty acids connected to glycerol are of the same type

e.g., tripalmitin.

Mixed triglycerides:

e.g., 1,3-bis(sn-3'-phosphatidyl)-sn-glycerol-2'-phosphate (phosphatidylcholine) and 1,3-bis(sn-3'-phosphatidyl)-sn-glycerol-2'-phosphate (phosphatidylcholine).

The main difference between fats and oils is for oils being liquid at room temperature, whereas, fats are solids.

This is mainly due to presence of larger percentage of unsaturated fatty acids in oils than fats that has mostly saturated fatty acids.

TAGs containing saturated fatty acids are solid at room temperature such as butter whereas

TAGs containing unsaturated fatty acids are liquid at room temperature such as olive oil. (Oleic acid, 18:1,9)

122 Lipid Chemistry-Triacylglycerols

Because the polar hydroxyls of glycerol and the polar carboxylates of the fatty acids are bound in ester linkages

Therefore the triacylglycerols are nonpolar, hydrophobic molecules, essentially insoluble in water

Triacylglycerols provide

Stored Energy and Insulation.

In vertebrates, specialized cells called adipocytes, or fat cells, store large amounts of triacylglycerols as fat droplets that nearly fill the cell

"

'''

Triacylglycerols are also stored as oils in the seeds of many types of plants.

Thus providing energy and biosynthetic precursors during seed germination

Adipocytes and germinating seeds contain lipases, enzymes that catalyze the hydrolysis of stored triacylglycerols.

Therefore releasing fatty acids for export to sites where they are required

123 Lipid Chemistry-Triacylglycerols (TAGs) Cont.

There are two significant advantages to using triacylglycerols as stored fuels, rather than polysaccharides such as glycogen and starch.

First, the carbon atoms of fatty acids are more reduced than those of sugars, and oxidation of triacylglycerols yields more than twice as much energy, as the oxidation of carbohydrates. Second, because triacylglycerols are hydrophobic and therefore unhydrated.

The organism that carries fat as fuel does not have to carry the extra weight of water of hydration that is associated with stored polysaccharides (2 g per gram of polysaccharide)

Moderately obese people with 15 to 20 kg of triacylglycerols deposited in their adipocytes could meet their energy needs for months by drawing on their fat stores

In contrast, the human body can store less than a day's energy supply in the form of glycogen (the polymer of glucose)

In some animals, triacylglycerols stored under the skin serve as insulation against low temperatures. Seals, penguins, bears and other warm-blooded polar animals are amply padded with triacylglycerols

124 Lipid Chemistry-Properties of TAGs

Physical properties

Neutral fats are

They are insoluble in water but soluble in organic fat solvents(e.g., ether, benzene, acetone, chloroform)

5. Specific gravity:

The specific gravity of all fats is less than 1.0, consequently all fats float in water

6. Emulsification:

Emulsions of fat may be made by shaking vigorously in water and by emulsifying agents such as gums and soaps

The emulsification of dietary fats in intestinal canal, brought about by bile salts, is a prerequisite for digestion and absorption of fats.

The bile salts, act to break apart the fat globules in the small intestines and allow them to become more "soluble" for absorption.

The hydrophobic fat molecules will clump together into globules in the watery mixture in the digestive system.

The emulsifiers break them down to smaller "globules" and allow them to become more soluble.

In Mustard variety of chemicals in the mucilage surrounding the seed hull act as emulsifiers

"

'''

125 Lipid Chemistry-Chemical Properties of TAGs

Chemical Properties of TAGs

Hydrolysis

The fats may be hydrolysed with super heated steam, by acids, or alkalies, by the specific fat splitting enzymes lipases to free fatty acids glycerol

Saponification

Hydrolysis of a fat by an alkali is called saponification

The resultant products are;
glycerol and

the alkali soaps

The number of mgs of NaOH/KOH required to saponify the free and combined FA in one gram of a given fat is called its saponification number

The amount of alkali needed to saponify a given quantity of fat will depend upon the number of carboxylic (COOH) group present

Thus fats containing short chain fatty acids will have more COOH groups per gram than long-chain fatty acids and this will result in a higher saponification number. And hence will have higher saponification number

Butter containing a larger proportion of short-chain fatty acids, such as butyric (C4) acid and caproic (C6) acid etc.

Therefore it has relatively high saponification number (195-205). In contrast, Olive Oil (which contains long-chain fatty acids) has saponification number of 195 or less.

Triolein is another example of simple TAG.

Tallow is the fat obtained from cattle used in soap industry.

126 Lipid Chemistry-Chemical Properties of TAGs Contd

2. Additive Reactions

The unsaturated fatty acids present in neutral fat exhibits all the additive reactions, i.e. hydrogenation, halogenation.

Hydrogenation

A hydrogenation reaction involves conversion of a carbon-carbon double bond to a carbon-carbon single bond through the addition of hydrogen

"

Hydrogenation As you continue to hydrogenate your molecule
Melting point increases

Fat becomes more solid at room temp

Hydrogenation

Oils which are liquid at ordinary room temperature, on hydrogenation become solidified

This is the basis of Banaspti ghee manufacturing.

Where inedible and cheap oils like cotton seed oil are hydrogenated and converted to edible solid fats.

The hydrogenation is done

under high pressure of hydrogen and is catalyzed by finely divided nickel or copper and
oleic acid of fats (liquid)

into stearic acid (solid).

unsaturated fats have lower melting points, stearic (SFA) melts at 70

127 Lipid Chemistry-Chemical Properties of TAGs Contd.

Halogenation

Similar to hydrogenation,

Halogens such as chlorine, bromine and iodine can also be added to double bonds in unsaturated fatty acids.

It is a very important property to determine the degree of unsaturation of the fat or oil that determines its biological value. 

The degree of unsaturation is reflected by Iodine number.

Iodine number is defined as the number of grams of iodine absorbed by 100 gm of fat.

The more the iodine number, the greater the degree of unsaturation.

Fats rich in saturated fatty acids have low iodine numbers,

while fats rich in unsaturated fatty acids have high iodine numbers

The determination of iodine number is useful to the chemist in determining the quality of an oil or its freedom from adulteration

Iodine number of cotton seed oil varies from 103 to 111.

That of olive oil from 79 to 88,

And that of linseed oil from 175 to 202

A commercial lot of olive oil which has iodine number higher than 88 might have been adulterated with cotton seed oil

The higher is the iodine number, the more reactive, less stable, more susceptible to oxidation and rancidification is the oil or fat.

128 Lipid Chemistry-Chemical Properties of TAGs Cont.

Chemical Properties (Contd.)

Rancidity

The chemical deterioration of fats.

When lipid-rich foods are exposed too long to the oxygen in air, they may spoil and become foul smelling.

Rancidity

Definition:

It is a physico-chemical change in the natural properties of the fat leading to the development of

unpleasant odor or

taste or

abnormal color

It occurs particularly

on aging

after exposure to atmospheric oxygen,

light,

moisture,

bacterial or fungal contamination

and/or heat.

Saturated fats resist rancidity more than unsaturated fats that have unsaturated double bonds.

Rancidity is due to

Hydrolytic rancidity: Formation of free fatty acids and glycerol

Oxidative rancidity: Formation of short chain aldehydes, ketones and dicarboxylic acids

which have objectionable taste and odor.

The unpleasant taste and smell associated with rancidity result from the oxidative cleavage of double bonds in unsaturated fatty acids

The oxygen of the air is necessary for this type of rancidity. This can be prevented by addition of anti-

oxidants such as vitamin E to foods.

II. Hydrolytic Rancidity

It is due to the slow hydrolysis of fats,

which in case of fats like butter results in the

liberation of short chain fatty acids which are volatile and have rancid taste and odor.

129

Lipid Chemistry-Waxes

Waxes

A second group of neutral lipids that are of physiological importance.

Although they are a minor component of biological systems.

Properties of waxes

Waxes are insoluble in water, but

soluble in fat solvents and are negative for acrolein test. very resistant to rancidity.

Waxes are not easily hydrolyzed as the fats and are indigestible by lipases

(enzymes responsible for hydrolysis of fats)

[r ≠ "r o - "k ° o "y p "x y "x ≠ ° s s y x k v "k v o " " " Waxes

are of two types:

[° ≠ o "Δ k - o • " " V ' r o ° Waxes or Non true waxes or

Wax-like compounds

True

Waxes are solid simple lipids containing a monohydric alcohol (with a higher molecular weight than glycerol

esterified to long-chain fatty acids.

"

'''

Triacontanol is a **fatty alcohol** of the general formula $C_{30}H_{62}O$, also known as **melissyl alcohol** or **myricyl alcohol**.

It is found in plant **cuticle waxes** and in **beeswax**.

"

Waxes are widely distributed in nature such as the secretion of certain insects as;

Bees-Δ k - 1 " " Z z o ° w k m ° s ' y p ' r o • z o ° w " Δ r k v " "

Waxe • "k v y " p ° w " z ° y ' o m s / o " n y k ' s q • " y p ' r o • u s • " k x n " p ° • " y p k x s v k v " k x n " " " leaves and fruits of plants.

Triacontanol is a fatty alcohol of the general formula $C_{30}H_{62}O$, also known as melissyl alcohol or myricyl alcohol.

It is found in plant cuticle waxes and in beeswax.

The name cetyl derives from the whale oil (Latin: cetus) from which it was first isolated

Cetyl Alcohol can be found in moisturizer, facial moisturizer, conditioner, antiaging, hair color, hair bleaching, facial cleanser. hand cream, shampoo, lipstick, eye cream.

130 Lipid Chemistry-True Waxes

1.True Waxes

Bees-wax is secreted by the honeybees that use it to form the combs. It

is a mixture of waxes

chief constituent is myricyl palmitate

(30C) (16C)

2.Spermaceti

is a wax that is most often found in the head cavities of the sperm whale.

Fatty esters are formed essentially of
cetyl palmitate and
cetyl myristate.

It was used in cosmetics, pharmacy and also in candles

recent international regulation concerning whale captures, has stopped its use.

2. Other Waxes or Non true
include esters of:

Cholesterol

Vitamin A

Vitamin D

and It
contains both free and esterified cholesterol, e.g., cholesterol-palmitate

Lanolin secretion helps sheep in reducing water evaporation from the skin.

It is used as industrial lubricant and in
cosmetics

131 Lipid Chemistry-Lipids and Nutrition

Dietary fats strongly influence the incidence of coronary heart disease (CHD).

In the past, dietary recommendations emphasized decreasing the total amount of fat in the diet.

Research now indicates that the type of fat is more important than the total amount of fat consumed

To gauge the effect of these dietary fats on CHD, we measure different biochemical parameters, which include:

LDL-Cholesterol

HDL-Cholesterol

Total Cholesterol

TAG

LDL and HDL stand for

Low Density Lipoprotein and

High Density Lipoprotein respectively

they are included in the class of complex lipids and serve to

transport lipids in the among others.

The risk increases progressively with higher values for serum total cholesterol and that of LDL cholesterol.

Association of TAGs is weaker than that of LDL cholesterol with CHD.

A much stronger correlation exists between the levels of blood LDL cholesterol and heart disease.

In contrast, high levels of HDL cholesterol have been associated with a decreased risk for heart disease

In order to understand the influence of these parameters on CHD, we must first understand biochemical role of Lipoproteins:

Lipoprotein particles are spherical aggregates with hydrophobic lipids at the core and hydrophilic protein side chains and lipid head groups at the surface.

These proteins are called apolipoproteins.

Lipoproteins function is to keep their component lipids soluble as they transport them in the aqueous environment of plasma.

Due to imbalanced metabolism of these lipoproteins there can be a gradual deposition of lipid especially cholesterol in tissues, which in arteries of the heart, can lead to CHD.

132 Lipid Chemistry-Lipoproteins

The lipoprotein particles include

chylomicrons (CM),

very-low-density lipoproteins (VLDL), low-density lipoproteins (LDL), and

high-density lipoproteins (HDL).

[rho = mass/volume] = density

• so 1" "no" = "kx" "• so" ypy" =

Different combinations of lipids and proteins produce particles of different densities ranging from chylomicrons to high-density lipoproteins

these particles can be separated by ultracentrifugation and HPLC

Chylomicrons are formed and assembled in intestinal mucosal cells after a fatty meal And carry dietary triacylglycerol, cholesterol, and cholesteryl esters to the peripheral tissues. As they are taken up by peripheral tissues in the form of fatty acids, the particle size decreases and density increases.

The remaining particle, called Chylomicron Remnant, is removed from the circulation by the liver.

133

Lipid Chemistry-Lipoproteins Contd

VLDLs are assembled in the liver. composed predominantly of TAGs synthesized in liver and contain some cholesterol and cholesteryl esters As VLDL pass through the circulation, TAG is degraded and taken up by peripheral tissues in the form of fatty acids, causing the VLDL to decrease in size and become denser, called VLDL remnant.

further removal of
 from these remnants produces low-density lipoprotein (LDL)
 LDL particles contain much less triacylglycerol than their VLDL predecessors,
 and have a high concentration of cholesterol and cholesteryl esters
 LDLs contain apoB-100 as their major apolipoprotein
 they carry cholesterol to extra hepatic tissues that have specific plasma membrane
 receptors that recognize apoB-100.
 apoB-100 receptors internalize the LDL into the cell.
 Remaining LDL is endocytosed by Liver cells.
 Oxidized LDL can also accumulate in the macrophage cells lining the arteries
 resulting in the formation of atherosclerosis.

134 Lipid Chemistry-Lipoproteins Cont.

HDL The fourth major lipoprotein type, high-density lipoprotein,

originates in the liver and small intestine
 as small, protein-rich particles that contain little cholesterol and no cholesteryl esters
 They take up cholesterol from non-hepatic (peripheral) tissues and return it to the liver as
 cholesteryl esters
 When cholesterol is taken up by HDL, it is immediately esterified and becomes
 hydrophobic and
 which is sequestered in the core of the HDL,
 This transforms the nascent HDL to a mature HDL particle.
 Mature HDL then returns to the liver, where the cholesterol is unloaded in a process called
 reverse cholesterol transport pathway

This is the basis for the inverse relationship seen between plasma HDL concentration and
 k' r o' y' m o' y' s 1' k x n' p' o' O K S' " n o' s x k' s y x' k' " r o' q y y n " m y o' o' y v n k' o' s' 3'

A much stronger correlation exists between the levels of blood LDL cholesterol and heart
 disease

In contrast, high levels of HDL cholesterol have been associated with a decreased risk for
 heart disease

Abnormal levels of plasma lipids (dyslipidemias) act in combination with smoking, obesity,
 sedentary lifestyle, insulin resistance, and other risk factors to increase the risk of CHD

Clinical studies have demonstrated that dietary or drug treatment of hyper -
 cholesterolemia is effective in decreasing LDL, increasing HDL, and reducing the risk for
 cardiovascular events.

135 Lipid Chemistry-

136 Lipid Chemistry-Role of Dietary Lipids

Role of dietary Lipids (Contd.)

Unhealthy Fat contains
 Trans fat
 Saturated Fat
 Increased cholesterol content

Trans fat

elevate serum LDL (but not HDL),

Therefore they increase the risk of CHD

Fatty acids of trans configuration in our food come from two different sources:

industrially produced partially hydrogenated fat (IP-TFA) and

ruminant produced Trans Fatty acid (RP-TFA)

Industrially produced trans fatt (IP-TFA)

These are made by partial hydrogenation of vegetable fat, and to a lesser extent, of fish oils by heating to about 400°C under high pressure and with the addition of different catalysts such as Nickel

Ruminant produced Trans Fatty acid (RP-TFA)

are made by bacterial metabolism of polyunsaturated fatty acids in the rumen of ruminants, such as cow & sheep

And are consequently present in all fats from these animals

The concentration of IP-TFA in partially hydrogenated fat may be as high as 60%.

Whereas the maximum content of RP-TFA in ruminant fat is about 6%.

In milk, RP-TFA is 4-6% of the fat.

Recent studies suggest that with equal amounts of intake

IP-TFA is more harmful than RP-TFA when compared on a gram-to-gram basis.

The deleterious effects of trans fats occur at intakes of 2 to 7 g/day

A single serving of French fries in a restaurant may contain this amount of trans fatty acids!

Therefore, to reduce the harmful effects of trans fats, one should

limit intake of foods prepared in IP-TFA containing oils such as baked goods and fast foods etc

3" "x• 'okn"yp"v s x q"p k "x'kuo"pyw"nks—• y ≠ °m• 3'

137 Lipid Chemistry-Role of Dietary Lipids Contd

Role of dietary Lipids (Contd.)

Saturated fats

Consumption of saturated fats is associated with high levels of

'y'kvz k•wk"m yv• 'o°yv'kxn"" "SKS"m yv• 'o°yv" "Zk'≠°k'on"p k'•"

l ≠ °o°1"" "r k'n"m oo•o• 1"" "Δryv"wsu1"" "kxsvkv'p k'• 1 "zkw"ys/l'kxn"" "nynyx≠'ys' 3""""

Among the SFAs, stearic acid (18:0) appears to have a neutral effect on LDL-C.

While lauric (12:0), myristic (14:0), and palmitic (16:0) acids are considered to be hypercholesterolemic

Saturated FAs increase plasma LDL-C by;

increasing the formation of LDL in tho'z k•wk"nywzk°'wox"" "kxn"l —"nomfok• x q"SKS" turnover.

The lowering of plasma LDL-C observed with PUFAs is likely due to;

redistribution of cholesterol between plasma and tissue pools

and up regulation of the LDL receptor

In contrast, monounsaturated fatty acids (MUFAs) and PUFAs of the (n-6) family have been shown to;

nonfok•o"zk•wk"mryv•'o'yv'myxmx'k'syx•"x"moxkv•'zn•"" " Dietary cholesterol:

Cholesterol is found only in animal products.

dietary cholesterol has little effect on plasma cholesterol.

Therefore, effect of dietary cholesterol on plasma cholesterol is less important than the amount and types of fatty acids consumed.

A further reduction in dietary cholesterol seems to be unnecessary in those people; who have already reduced their intake of saturated fat

and increased the ratio of polyunsaturated to saturated fatty acids

The Mediterranean Diet

Mediterranean cultures, show a low incidence of coronary heart disease

The Mediterranean diet is an example ypk"" "ns""m"x"wyxy#x•k'°k'on"pk''-kms•"

-pyw"yso"ys"xkn" -3 fatty acids (from fish oils and some nuts),

but low in saturated fat

The Mediterranean diet contains

seasonally fresh food, with

an abundance of plant material,

low amounts of red meat, and

olive oil as the principal source of fat

The Mediterranean diet is associated with decreased serum total cholesterol

and LDL cholesterol but

little change in HDL cholesterol when compared with a typical Western diet higher in saturated fats

138 Lipid Chemistry-Structure of Phospholipids

STRUCTURE OF PHOSPHOLIPIDS

There are two classes of phospholipids:

those tr k"rk\o"qvno°yvk•"k"l kmul yxoAqvno°yzry•zryvzn•"" " those that contain sphingosine:

sphingophospholipids

Phospholipids are the predominant lipids of cell membranes

Membrane lipids are amphipathic i.e. one end of the molecule is hydrophobic, the other hydrophilic

Their hydrophobic interactions with each other and their hydrophilic interactions with water direct their assembly into bilayers. The formation of membrane bilayers

help in partitioning the cellular environment from extracellular environment.

Non-membrane-bound phospholipids serve additional functions, for example, as components of lung surfactant

essential components of bile. 

In contrast to triacylglycerol which is essentially synthesized only in liver, adipose tissue, lactating mammary glands, and intestinal mucosal cells essentially all cells except mature erythrocytes can synthesize phospholipids.

139 Lipid Chemistry-Glycerophospholipids

Glycerophospholipids

Phosphatidic acid

It is the simplest phosphoglyceride,

It is a diacylglycerol with a phosphate group on the third carbon of glycerol.

Phosphatidic acid is the precursor of the other members of this group.
 Further esterification with a low-molecular weight alcohol gives a glycerophospholipid
 In general, glycerophospholipids contain
 a C16 or C18 saturated fatty acid at C-1 and
 a C18 or C20 unsaturated fatty acid at C-2 and a phosphate group on C-3

The phosphate group on phosphatidic acid (PA) can be esterified to another compound containing a hydroxyl group

phosphatidylserine
 phosphatidylethanolamine
 phosphatidylcholine

The
a
The

molecule is both a
primary amine and
primary alcohol
fatty acids in

glycerophospholipids can be any of a wide variety, so a given phospholipid (phosphatidylcholine, for example) may consist of several molecular species, each with its unique complement of fatty acids.

For instance, Dipalmitoyl phosphatidyl choline and Palmitoyl-oleylphosphatidylcholine are two different examples of Lecithins.

140 Lipid Chemistry-Phosphotidylcholines

Glycerophospholipids

(Contd.)

Phosphatidylcholines (Lecithins)

the most abundant phospholipids of the cell membrane represent a large proportion of the body's store of choline- important in nervous transmission, as acetylcholine.

Dipalmitoyl phosphatidyl choline (DPPC or dipalmitoylecithin), is also the major lipid component of lung surfactant

Alveoli are the structural and functional unit of respiratory system in which gaseous exchange takes place.

A thin fluid layer lines the alveoli for efficient gas exchange.

When the water forms a surface with air, the water molecules on the surface (in contact of air) are strongly attracted to each other trying to reduce the surface area of contact.

This is called "surface tension".

surface tension might result in alveolar collapse

it requires a certain inflation pressure to maintain expanded alveoli

the higher the surface tension, the more pressure required to inflate the bubble, especially in small alveoli.

made and secreted by lung cells, surfactant serves to decrease the surface tension of this fluid layer.

It scatters among the fluid molecule decreasing the attraction between them.

surfactant have both hydrophilic and hydrophobic regions

by adsorbing to the air-water interface of alveoli hydrophilic head groups in the fluid and the hydrophobic tails facing towards the air (DPPC) reduces surface tension.

141 Lipid Chemistry-Phosphotidyl Ehanol Amine

Glycerophospholipids (Contd.)

Phosphatidyl ethanol amine and phosphatidylserine are also found in cell membranes and differ from phosphatidylcholine only in that ethanolamine or serine, respectively, replaces choline
Serine is a standard amino acid which on decarboxylation produces ethanolamine: a primary alcohol and a primary amine.

Phosphatidylserine also plays a role in apoptosis (programmed cell death)
Cardiolipin

diphosphatidylglycerol

Two molecules of phosphatidic acid esterified through their phosphate groups to an additional molecule of glycerol

In eukaryotes, cardiolipin is virtually exclusive to the inner mitochondrial membrane, where it appears to be required for the maintenance of certain respiratory complexes of the electron transport chain.

Decreased cardiolipin levels or alterations in its structure or metabolism cause mitochondrial dysfunction in aging and in pathological conditions including heart failure

Cardiolipin is antigenic, and is recognized by antibodies raised against *Treponema pallidum*, the bacterium that causes syphilis

The VDRL test measures immunoglobulin G (IgG) and IgM antibodies to lipoidal material released from damaged host cells as well as to lipoprotein-like material and possibly cardiolipin released from the treponemes

Anti-cardiolipin antibodies may also be found non-specifically in certain autoimmune diseases.

142 Lipid Chemistry-Phosphatidylinositol Glycerophospholipids (Contd.)

Phosphatidylinositol

consists of phosphatidic acid and Inositol in an ester linkage

Inositol is a polyol synthesized in the body from glucose and have the same chemical formula as glucose

Phosphatidylinositol (PI) is an unusual phospholipid in that it often contains stearic acid on carbon 1 and arachidonic acid on carbon 2 of the glycerol

PI, therefore, as a reservoir of arachidonic acid, serves as precursor for prostaglandin synthesis

Specific proteins can be covalently attached via a carbohydrate bridge to membrane-bound PI.

Second messengers

The phosphorylation of membrane-bound phosphatidylinositol produces phosphatidylinositol 4,5-bisphosphate (PIP₂)

The degradation of PIP₂ by phospholipase C occurs in response to the binding of a variety of neurotransmitters, hormones, and growth factors to receptors on the cell membrane

The products of this degradation are inositol 1,4,5 trisphosphate (IP₃) and diacylglycerol (DAG)

These products mediate the mobilization of intracellular calcium and the activation of protein kinase C, respectively, which act synergistically to evoke specific cellular responses

143 Lipid Chemistry-Ether Lipids

Ether lipids

They are type of Glycerophospholipids, in which one acyl chains is attached to glycerol in ether linkage, rather than ester linkage. Ether lipids with an unsaturated group (alkenyl) at the 1st position on the glycerol chain are called Plasmalogens

||||

|||

"

In some instances serine, or inositol may be substituted for ethanolamine or choline

These compounds constitute as much as 10% of the phospholipids of brain and muscle

Vertebrate heart tissue is uniquely enriched in ether lipids

About half of the heart phospholipids are plasmalogens

The membranes of halophilic bacteria, ciliated protists, and certain invertebrates also contain high proportions of ether lipids.

The functional significance of ether lipids in these membranes is unknown; perhaps their resistance to the phospholipases that cleave ester-linked fatty acids from membrane lipids is important in some roles

Platelet-activating factor (PAF)

This is an unusual ether glycerophospholipid, with a saturated alkyl group in an ether link to carbon 1 and an acetyl residue (rather than a fatty acid) at carbon 2 of the glycerol backbone

PAF is synthesized and released by a variety of cell types
It binds to surface receptors, triggering potent thrombotic and acute inflammatory events
It causes platelets to aggregate and degranulate (required for clotting), and neutrophils and alveolar macrophages to generate superoxide radicals (required for microbial killing)

144 Lipid Chemistry-Sphingolipids

SPHINGOLIPIDS

Sphingolipids, like other membrane lipids, are composed of a
hydrophobic portion, (ceramide) and
a polar head group

The first three carbons at the polar end of sphingosine are analogous to the three carbons of glycerol in glycerophospholipids

The amino group at C-2 is usually protonated. The fatty acid is usually saturated or monounsaturated, with 16, 18, 22, or 24 carbon atoms.

Ceramide is the parent compound for this group

SPHINGOLIPIDS

There are two subclasses of sphingolipids:

Sphingomyelins

Sphingoglycolipids

Sphingomyelins

contain phosphocholine or phosphoethanolamines as their polar head group and are therefore classified along with glycerophospholipids as phospholipids

Sphingoglycolipids

are molecules that contain both
carbohydrate and
lipid (in the form of ceramide) components

"

When the alcohol group at carbon 1 of sphingosine is, esterified to **phosphorylcholine**, **sphingomyelin**, the only significant sphingophospholipid in humans, is produced

Sphingomyelins are

present in the plasma membranes of animal cells and are especially prominent in nerve tissue including myelin, -thus the name "sphingomyelins

Sphingomyelin of the **myelin sheath** contains predominant longer-chain fatty acids such as lignoceric acid and nervonic acid (24 carbon)

whereas **gray matter of the brain** has sphingomyelin that contains primarily stearic acid(18 carbon)

"

145 Lipid Chemistry-Glycosphingolipids

Glycosphingolipids are molecules that contain both

carbohydrate and lipid components

Like the phospholipid sphingomyelin, they are derivatives of ceramide

They are also an important component of membrane bilayers.

They occur largely in the outer face of plasma membranes have polar head groups with one or more sugars connected directly to the -OH at C-1 of the ceramide moiety by an O-glycosidic bond

Cerebrosides

have a single sugar linked to ceramide;

galactocerebrosides are found in the plasma membranes of neurons

glucocerebrosides in the plasma membranes of non neural cells

Galactosylceramide is a major glycosphingolipid of brain and other nervous tissue,
It contains a number of characteristic C24 fatty acids,
eg, cerebronic acid

Globosides

are glycosphingolipids

with two or more sugars, usually

D -glucose,

D- galactose, or

N-acetyl-D-galactosamine.

A glycosphingolipid that has only one sugar as the side chain is called a cerebroside

Globosides are highly abundant in RBCs.

Globosides are also found in human serum, spleen and liver.

"

146 Lipid Chemistry-Glycosphingolipids Contd

Gangliosides

They have oligosaccharides as their polar head groups and one or more residues of N-acetylneuraminic acid (a sialic acid), at the termini.

D -glucose,

D- galactose, or

N-acetyl-D-glucosamine-6-phosphate " N-acetylneuraminic acid

Sialic acids are acidic sugars with a nine-carbon backbone, of which the most common is N-acetylneuraminic acid

Sialic acid gives gangliosides the negative charge at pH 7

Gangliosides with one sialic acid residue are in the GM (M for mono-) series, those with two are in the GD (D for di-) series, and so on (GT three sialic acid residues; GQ, four)

About 6% of brain lipids are gangliosides and were first isolated from the ganglion of brain cells

They act as specific receptors for glycoprotein hormones in the cells.

Some gangliosides also serve as receptors for some bacterial protein toxins e.g. Cholera toxin binds to the GM1 gangliosides on the surface of target cells

Gangliosides also help in cell-cell recognition and thus have a significant role in growth and differentiation of tissues and also in carcinogenesis

Sulfatides

Sulfoglycosphingolipids (sulfatides) are cerebroside that contain sulfated galactosyl residues, and are therefore negatively charged at physiologic pH

"Sulfatides are found predominantly in nerve tissue and kidney"

147 Lipid Chemistry-ABO Blood Groups and Glycosphingolipids

ABO Blood Groups and Glycosphingolipids

The basis of blood groups depend on expression of different antigens on RBCs.

These antigens are found in cell membranes as oligosaccharide components of glycosphingolipids
glycoproteins

Different compositions of carbohydrate (oligosaccharide chains) in different antigens. These oligosaccharides are attached to proteins through a serine or threonine residue or to ceramide lipid intermediate

The precursor to the ABO blood group antigens, present in people of all common blood

types, is the O antigen (also referred to as H antigen.)

Modification of this O antigen then

results in A, B or AB blood groups respectively.

'''

"

These antigens are present on other cells of the body as well.

148 Lipid Chemistry-Glycerophospholipids

Glyceroglycolipids

These are the predominant membrane lipids of the plants, such as those of chloroplast. They include Galactolipids and

Sulfolipids

As the name indicates these glycolipids contain glycerol instead of sphingosine as their backbone.

Galactolipids

In galactolipids, the C3 of the glycerol moiety is connected to one or more galactose residues by glycosidic linkages.

The head groups of these galactolipids are uncharged but polar.

Galactolipids are localized in the internal membranes of chloroplasts. They constitute about 70% to 80 % of plant membrane lipids and thus are the most abundant lipids in the biosphere.

Sulfolipids

They are membrane glycolipids with sulfur containing functional

groups. They are connected to the glycerol backbone by glycosidic linkage

149 Lipid Chemistry-Steroids and Cholesterol

Steroids and Cholesterol

A steroid is a lipid whose structure is based on the tetracyclic (four-ring) structure consists of

3 cyclohexane rings.

1 cyclopentane ring.

Steroids with eight to ten carbon atoms in the side chain at C-17 and a hydroxyl group at C-3 are classified as sterols

Cholesterol is the major sterol in animal tissues

Cholesterol has an eight carbon branched hydrocarbon chain attached to C-17 of the steroid nucleus. Ring A has a hydroxyl group at C-3, and ring B has a double bond between C-5 and C-6

Cholesterol is an amphipathic lipid

Cholesterol is a structural component of all cell membranes, modulating their fluidity

Cholesterol is a precursor of bile acids, steroid hormones, vitamin D

All tissues containing nucleated cells are capable of cholesterol synthesis, which occurs in the endoplasmic reticulum and the cytosol

Cholesterol is present in tissues and in plasma either as free cholesterol or combined with a long-chain fatty acid as cholesteryl ester

In plasma, both forms are transported in lipoproteins

in esterified form, with a FA attached at C-17. The free cholesterol is found in the plasma and in the cell membranes.

Cholesteryl esters are not found in membranes, and are normally present only in low levels in most cells

The ring structure of cholesterol cannot be metabolized to CO₂ and H₂O in humans

Therefore humans cannot utilize cholesterol for energy.

Cholesterol is excreted from the body via the bile either in the unesterified form or after conversion into bile acids in the liver

150 Lipid BILE ACIDS AND BILE SALTS

Bile helps with lipid digestion

Bile can either pass directly from the liver into the duodenum, or be stored in the gallbladder when not immediately needed for digestion

Bile consists of a watery mixture of organic and inorganic compounds

Phosphatidylcholine (lecithin) and

bile salts (conjugated bile acids) are quantitatively the most important organic components of bile

The primary bile acids are synthesized in the liver from cholesterol

chenodeoxycholic acid

'''

"

The bile acids contain 24 carbons, with two or three hydroxyl groups and a side chain that terminates in a carboxyl group
The carboxyl group has a pKa of about 6 and, is not fully ionized at physiologic pH- hence the term "bile acid"
The bile acids are amphipathic molecules
They therefore can act as emulsifying agents in the intestine

They emulsify dietary triacylglycerol and other cywzv-''' " for degradation by pancreatic digestive enzymes.

151 Lipid Chemistry-Eicosanoids

Eicosanoids are a large group of lipid messengers with potent effects on every tissue in the body

Eicosanoids are derived from metabolism of 20-carbon, polyunsaturated fatty acids

Prostanoids consisting of

Prostaglandins

Prostacyclins

Thromboxanes

1. Leukotrienes

2. Lipoxins

3. Epoxides

These extremely potent compounds acting through their specific receptors elicit a wide range of physiologic and pathologic responses.

particularly important in eliciting inflammatory response that occurs after infection or injury and

produce symptoms such as pain, swelling, and fever.

they also control bleeding through forming blood clots

Eicosanoids are derived from either omega-3 or omega-6 fatty acids.

Arachidonic acid is the most common precursor of the eicosanoids.

Three major pathways for the metabolism of arachidonic have been discovered so far.

Cyclooxygenase pathway; prostaglandins and thromboxanes.

lipoxygenase pathway; leukotrienes, HETEs, and lipoxins.

cytochrome P450 pathway epoxides and HETEs

There are different series of eicosanoids

depending on the precursor 20 C FA. derived from the essential fatty acids linoleic acid and linolenic, OR

directly from dietary arachidonic acid and eicosapentaenoic acid

Depending on the precursor different numbers of double bonds are present in these eicosanoids reflecting the parent 20C FA.

152 Lipid Chemistry- Cyclooxygenase Pathway

Prostaglandins

Wf•'kqkxns•"k°"p''-kn•"nyx'ksxq"75"nk°l yx"kyw•1xmwnsq"" "kx"so°xkv: - carbon ring.

Prostaglandins have
a hydroxyl group at C 15,
a double bond between C 13 and C 14, and
various substituents on 5 membered ring at C9 and C11

Nomenclature of prostaglandins (PGs) involves the
k••sqxwox"ypk"nkzskv'o'o°"-WNL.1"" "k"x#w°kv°≠l •miz'"-WNL6.1kxn""
for the PGF faw s-1k"N°ou"v'o'o°"•≠l •miz'"-WNM7 .3""
The capital letter refers to the ring substituents at positions X and Y.

The numeral subscript that follows the capital letter(e.g. PGE1) refers to the PG series
1, 2, or 3, determined by the number of unsaturated bonds present in the linear portion
of the hydrocarbon chain.

It does not include double bonds in the internal ring.
The double bonds between carbons 13 and 14 are trans; the others are cis
The Greek letter subscript, found only in the F series, refers to the position of the
hydroxyl group at carbon 9.
[r 9 "r -n°y—vq°y#z"z°svk°s—"o—9 ° "x"r o" "zosition, i.e. it lies below the plane of the
ring.
Prostaglandins have a hydroxyl group at C 15, a double bond between C 13 and C 14,
and various substituents on 5 membered ring at C9 and C11 Double bonds also may
be present between carbons 5 and 6 and between carbons 17 and 18 in case of series
2 and 3

153 Lipid Chemistry-Cyclooxygenase Pathway (Contd.)

CYCLOOXYGENASE PATHWAY (Contd.)

Thromboxanes (TX)
also formed via the cyclooxygenase pathway,
differ from PGs in that they contain a 6-membered ring that includes an oxygen atom

Biosynthesis of the PGS and TX

Those derived from arachidonic acid, the 2-series, (such as PGE₂, TXA₂), are described here. The 1-series and the 6-series and the 3-series are present in very small amounts in humans.

The initial step, which is catalyzed by a cyclooxygenase (COX), forms the five-membered ring and adds four atoms of oxygen (two between C 9 and C11, and two at C 15) to form the unstable PGG₂.

The hydro-peroxy group at carbon 15 is

~#smv"on#non"y"k"r-n°y—vq°y#z"l —k"zo°y—sk•o"y"p°w"WO7" " PGH₂ is the precursor of all other PGs and TXAs

The next step is tissue specific

For example, PGH₂ may be reduced to PGE₂ or PGD₂ by specific isomerases (PGE synthase or PGD synthase respectively)

3" "My"o—kwzw1[bH"Z—x'r k•o"s "z°o•ox""x'r sgr "nyxn0x°k'syx"x'z'k'ov'•"kxn"p°w•" TXA₂.

In the vascular endothelium, however, PGH₂ is converted to the PGI₂ by action of Prostacyclin Synthase.

The beneficial effect of cold water fish (e.g., salmon), with a high content of eicosapentaenoic (EPA), and docosa-hexaenoic acid (DHA) comes from the fact that they lead to formation of more TXA₃ relative to TXA₂.

TXA₃ is less effective in stimulating platelet aggregation than its counterpart in the 2-series, TXA₂.

Platelet aggregation is the culminating step in the cardiovascular diseases due to atherosclerosis.

154 Lipid Chemistry-Lipoxygenase Pathway

Lipoxygenase Pathway:

Synthesis of the Leukotrienes, and Lipoxins (Count.)

In addition to serving as a substrate for the cyclooxygenase pathway, arachidonic acid and other 20 C FAs also act as substrate for the lipoxygenase pathway.

In contrast to products of the cyclooxygenase pathway which are cyclical.

Products of lipoxygenase pathway are linear

The series starts from 3.

the eicosanoids synthesized in lipoxygenase pathway from;

Eicosa-tri-enoic acid (ETA) have 3 double bonds

from; Arachidonic acid have 4 double bonds

From; Eicosa-pentaenoic acid (EPA) have 5 double bonds

3" "[r o"vzy—qoxk•o"ox" —wo•"nk'kv—o"r o"smγ°zy°k'syx"yp'kn oxygen molecule on to a carbon of one of several double bonds of arachidonic acid, forming a hydro-peroxy (OOH) group at these positions

The double bonds at which oxygen is added are between C5 & C6, between C11 and C12 and between C14 and C15.

155 Lipid Chemistry-Lipoxygenase Pathway (Contd)

Lipoxygenase Pathway:

Synthesis of the Leukotrienes, and Lipoxins (Contd.)

When the oxygen is added, to the double bond it isomerizes to a position one carbon away from the hydro-peroxy group and is transformed from the *cis* to the *trans* configuration

As a result of this isomerization, double bond

between C5 & C6 moves to C6 & C7,

between C11 & C12 moves to C10 & C11

between C14 & C15 moves to C13 & C14

The hydro-peroxy group is unstable and

can be converted to the more stable hydroxy group 'y"p^w"OL[L•" " HETEs themselves act as messenger molecules

The other alternate is the conversion of HPETEs to leukotrienes(LTs) and Lipoxins(LXs),
Which are more potent and have more defined physiological roles.
The leukotrienes(LTs) and Lipoxins(LXs),
Have 3 to 5 double bonds as described earlier, in comparison to
prostaglandins which have 1 to 3 double bonds.

156 Lipid Chemistry-Lipoxygenase Pathway (Contd.)

Lipoxygenase Pathway:

HPETEs (Hydro-per-oxy-eicosa-tetra-enoic acids) are the precursors of leukotrienes and lipoxins

The major leukotrienes are produced by 5-lipoxygenase. 5-HPETE is converted to leukotriene A₄ (LTA₄).

Other functional leukotrienes are formed from LTA₄. LTA₄ is converted to LTB₄, as a 5,12-dihydroxy derivative.
the addition of reduced glutathione to carbon 6 forms LTC₄

LTD₄ on removal of glycine becomes LTE₄

Leukotrienes were so named because they were first discovered in leukocytes(white blood cells)

157 Lipid Chemistry-Lipoxygenase Pathway (Cont)

Synthesis of the Leukotrienes, and Lipoxins (Contd.)

The lipoxins are formed through the action of 15-lipoxygenase followed by the action of 5-lipoxygenase on arachidonic acid.

A series of reductions of the resultant hydro-per-oxy groups leads to the formation of tri-hydroxy derivatives of arachidonic acid known as the lipoxins.

Lipoxins induce chemotaxis and stimulate superoxide radicals for killing of microorganisms

Prostaglandins, thromboxanes, leukotrienes and lipoxins have very short half lives and rapidly degraded in the body.

In summary, Eicosanoids are derived from C20 (eicosanoic) fatty acids synthesized from the essential fatty acids and make up important groups of physiologically active compounds.

158 Nucleotides and Nucleic Acids

Biomedical importance of nucleotides

Precursors of nucleic acids: Nucleotides are the building blocks of nucleic acids.

Without them, DNA or RNA can not be produced.

Transmission of genetic information;

This gives them the ability to store and transmit genetic information from;
one generation to the next which is a fundamental condition for life

Protein synthesis,

the ultimate expression of this information, is therefore dependent on nucleotides.

Energy currency: Nucleotides play an important role as "energy currency" in the cell.
Nucleoside tri- and diphosphates such as ATP and ADP are the principal donors and
acceptors of phosphoryl group in metabolism.

By doing this, they play a key role in the energy transduction.

This energy is used in almost every energy requiring process in the body, such as;
Muscle contraction, Transmission of nerve impulse, Transports of nutrients across cell
membrane Motility of spermatozoa And many more energy dependent processes

159 Nucleotides and Nucleic Acids- Biomedical importance of nucleotides

Carriers of intermediates: Nucleotides also serve as carriers of activated intermediates in the
synthesis of some carbohydrates, lipids, and proteins.

The sugar derivatives UDP-glucose and UDP-galactose participate in sugar inter
conversions

And in the biosynthesis of starch and glycogen

Similarly, nucleoside-lipid derivatives such as CDP acylglycerol are intermediates in
lipid biosynthesis.

Co-enzymes:

When linked to vitamins nucleotides are structural components of several essential coenzymes, for example, coenzyme A,

FAD, (Flavin Adenine Dinucleotide) and Co-enzymes:

NAD⁺ (Nicotinamide adenine Dinucleotide) and

NADP⁺ (Nicotinamide adenine Dinucleotide Phosphate)

Regulatory compounds: Nucleotides are important regulatory compounds for many of the pathways of intermediary metabolism, inhibiting or activating key enzymes.

Roles that nucleotides perform in metabolic regulation include:

ATP-dependent enzyme phosphorylation in key metabolic reactions.

Allosteric regulation of enzymes by ATP, AMP, and CTP

Phosphorylation can either activate or inhibit ATP- dependent enzyme depending on the type of enzyme.

Similar is the case with dephosphorylation.

160 Nucleotides and Nucleic Acids- Biomedical importance of nucleotides (Contd)

Sulfate group donor: Adenosine 3'-phosphate-5'-phosphosulfate is the sulfate donor for
• synthesis of sulfate conjugates of drugs.

Methyl group donor:

S-adenosylmethionine (SAM) is the methyl group donor for
e.g Nor-adrenaline Adrenaline by methylation.

Nucleotides, such as cyclic AMP (cAMP) and cyclic GMP (cGMP),

act as second messengers in signal transduction pathways.

Signal Transduction: GTP and GDP play key roles in activating or inhibiting proteins in various cellular signaling cascades.

Thymine, uracil, and cytosine are purine and pyrimidine analogs that contain halogens, thiols, or additional nitrogen atoms;

Their uses include chemotherapy for cancer

as suppressors of the immune response during organ transplantation.

as anti-viral drugs such as in the treatment of AIDS

161 Nucleotides and Nucleic Acids-Composition of Nucleotides

Nucleotides are composed of

A nitrogenous base (purine or pyrimidine)

A pentose monosaccharide

One, two, or three phosphate groups

Nitrogenous Bases

The nitrogen-containing bases belong to two families of compounds:

Purines

Pyrimidines

By the attachment of different groups to the rings, different types of pyrimidine and purine are generated.



However, there are some exceptions such as naming of uracil

The utility of these nitrogen-containing ring structures lies in the ability of the nitrogen to

form hydrogen bonds and to

accept and donate electrons while still part of the ring.

162 Nucleotides and Nucleic Acids- Composition of Nucleotides (Contd.)

Purines

Both DNA and RNA contain the same purine bases:

Adenine (A)

Guanine (G)

Adenine when combines with pentose the structure is known as Adenosine or deoxyadenosine

Guanine when combines with pentose the structure is known as Guanosine or deoxyguanosine


Adenine is 6-aminopurine

Guanine is 2-amino,6-hydroxypurine

Minor Purine Bases: Inosine (I) & methyl guanine (7mG)

Unnatural: Mercaptopurine, Allopurinol & 8-Azaguanine

Other purines include: hypoxanthine and xanthine



Uric acid is the catabolic end product of purines in human beings.

163 Nucleotides and Nucleic Acids- Composition of Nucleotides (Cont.)

Pyrimidines:

Pyrimidines include: "J —y•xo"-J . in both DNA and RNA

Thymine (T) only in DNA

Uracil (U) only in RNA

Cytosine when combines with pentose it becomes deoxycytidine and cytidine
 Thymine becomes thymidine and deoxythymidine
 Uracil (U) becomes uridine and deoxyuridine depending on the type of sugar.
 Cytosine is 2-oxy-4-amino-pyrimidine
 Thymine is 2,4-dioxy-5-methyl-pyrimidine " Uracil is 2,4-dioxy-pyrimidine

T and U differ by only one methyl group, which is present on T but absent on U

Minor Pyrimidine Bases: "Ks -n°y°nso"-KO\ ."1'

5-T o'r -vJ -'knsx°+"'" ": -Hydroxy-Methyl Cytadine

Unnatural Pyrimidine Bases:

Fluorouracil (5FU) &

6-Aza Cytosine (AZC)

Unlike the purine ring, which is not cleaved in human cells, the pyrimidine ring is opened and degraded

'y"r q'r v" yv'l v"z°yn°m° 1' -k°kxso°kxn" -amino-iso-butyrate,
 with the production of NH₃ and CO₂.

164 Nucleotides and Nucleic Acids- Composition of Nucleotides (Contd)

Pentose Sugar

D-ribose and 2-deoxy D-ribose are the only sugars so far found in the nucleic acids.



These also pentoses belong to D-family

[r o—"k°o"z°o°ox "'k° "M°kxy°o"-°xq."x"'r o"p°w"yp° -Anomer

The addition of a pentose sugar to a base produces a nucleoside
If the sugar is D-ribose, a ribonucleoside is produced
If the sugar is 2-deoxy D- ribose, a deoxyribonucleoside is produced

Phosphate group

There may be one, two, or three phosphate groups present in nucleotides.
Nucleotides are monophosphate, diphosphate, or triphosphate esters of nucleosides

These phosphate groups give an over all negative charge to the nucleotides.

165 Nucleotides and Nucleic Acids- Properties of Nitrogenous Bases

Properties of Nitrogenous Bases

Aromatic: The Nitrogen containing bases are aromatic i.e. they have alternate double

lyxn• "" **Heterocyclic:**

They are heterocyclic i.e. structures that contain other atoms in addition to carbon, such as nitrogen in the ring structure

The six-atom rings of purines and pyrimidines are numbered in opposite directions.

Weak Bases: Purines or pyrimidines with an NH_2 group are weak bases

Functional Groups: The most important functional groups of pyrimidines and purines are
ring nitrogens
carbonyl groups
exocyclic amino groups

Hydrophobicity: They are insoluble in water at the near-neutral cell pH

Stacking Interaction: Hydrophobic stacking interactions in which two or more bases are positioned with the planes of their rings parallel (like a stack of coins) are one of two important modes of interaction between bases in nucleic acids.

Base stacking helps to minimize contact of the bases with water, and these interactions are very important in stabilizing the three-dimensional structure of nucleic acids.

166 Nucleotides and Nucleic Acids- Properties of Nitrogenous Bases (Contd)

UV light absorbance: The conjugated double bonds of purine and pyrimidine derivatives absorb ultraviolet light.

Nucleic acids are characterized by a strong absorption at wavelengths near 260 nm. The mutagenic effect of ultraviolet light is due to its absorption by nucleotides that results in chemical modifications in DNA.

This property is also utilized in quantitative and qualitative analysis of nucleotides and nucleic acids.

Tautomerism: All these bases can exist in keto-enol or amine-imine form. At physiologic pH keto and amine form is predominant.

Note that the smaller pyrimidine molecule has the longer name and " the larger purine molecule the shorter name.

167 Nucleotides and Nucleic Acids- N-glycosidic bond

N-glycosidic bond

Sugars are linked to the nitrogenous base by an N-glycosidic bond, almost always to the N-1 of a pyrimidine
N-9 of a purine

The N-glycosyl bond is formed by removal of the elements of water
a hydroxyl group from the pentose and
hydrogen from the base

Thus it is a condensation reaction.

Similar to O-glycosidic bond formation in carbohydrates

However, N-glycosidic bonds, have Nitrogen atom instead of oxygen linking the two
°o•n#o•3" "r o"knnsyx"yp"r o"qv•n•nll yxn"y"xsy°qoxous base is indicated by the
name change

such as from adenine to adenosine for the glycosidic bond

168 Nucleotides and Nucleic Acids-Numbering of Carbon and Nitrogen Atoms

Numbering of Carbon and Nitrogen Atoms

The carbon and nitrogen atoms in the rings of the base and the sugar are numbered separately

3" "l r o"k'ýw•"x"r o"rings of the bases are numbered

1 to 6 in pyrimidines &

1 to 9 in purines

In the pentoses of nucleotides and nucleosides the carbon numbers are given a prime (') designation to distinguish them from the numbered atoms of the nitrogenous base.

The carbons in the pentose are numbered 1' to 5 '.

Numerals with a prime (e.g., 2' or 3') distinguish atoms of the sugar from those of the heterocycle.



Thus, when the 5'-carbon of a nucleoside (or nucleotide) is referred to, a carbon atom in the pentose, rather than an atom in the base, is being specified.

169 Nucleotides and Nucleic Acids- Phosphodiester Bond

Phosphodiester Bond

When two or more nucleotides combine together a phosphodiester bond is formed.

The 5' carbon of one sugar is linked to the 3' carbon of another sugar by a phosphodiester bond. This bond is formed by the reaction of the 5' phosphate group of one sugar and the 3' hydroxyl group of another sugar, releasing a water molecule.

By definition, the 5' end lacks a nucleotide at the 3' position. The 3' end lacks a phosphate group.

The sugar and phosphate group is called the backbone of the nucleic acid.

The backbones of both DNA and RNA are hydrophilic.

The hydroxyl groups of the sugar residues form hydrogen bonds with water.

The phosphate groups, are completely ionized and negatively charged at pH 7
 The negative charges are generally neutralized by ionic interactions with positive charges which are present on proteins, metal ions, and polyamines.
 Depending on the number of nucleotides
 mono- di-, tri-, oligo- and polynucleotides are formed
 as a result of phosphodiester bond formation.

170 Nucleotides and Nucleic Acids-Cyclic Nucleotides

Cyclic Nucleotides

There are two important cyclic nucleotides:

Cyclic AMP cAMP

Cyclic GMP cGMP

Cyclic AMP is a cyclic nucleotide

cAMP is $\text{3',5'-bisphosphate}$ chemically it is 3'-5' adenosine monophosphate.

Functions of c-AMP

Acts as second messenger in the cell

It has role in glycogen metabolism

cAMP, glycogenolysis

cAMP TAG metabolism

cAMP lipolysis

It decreases cholesterol synthesis

It causes activation of protein kinases which in turn;
 activate or deactivate other enzymes.

It regulates the cell membrane permeability, by increasing permeability of cell membrane

to H_2O , Na^+ , K^+ & Ca^{+2}

Moreover, it regulates

insulin secretion,

catecholamine biosynthesis & Melatonin synthesis

Cyclic GMP is synthesized from GTP

It serves as a second messenger in response to nitric oxide during relaxation of smooth muscle (especially blood vessels) so it has role in smooth muscle relaxation and vasodilatation.

It also has role in

Protein phosphorylation

Neurotransmission

Insulin action

Regulation of sodium channels

DNA

DNA is present in nuclear chromosomes of eukaryotes

DNA is a polymer of deoxyribonucleoside monophosphates

DNA is a repository of genetic information

It is found associated with basic proteins HISTONES and also various other proteins present in nucleus (nucleoproteins)

The complex folding of large chromosomes within eukaryotic chromatin and bacterial nucleoids is generally considered tertiary structure

The most important functional groups of pyrimidines and purines are:

"

Hydrogen bonds involving the amino and carbonyl groups are the second important mode of interaction between bases in nucleic acid molecules.

The most important hydrogen-bonding patterns are those defined by James D. Watson


These two types of base pairs predominate in double-stranded DNA and RNA.

173 Nucleotides and Nucleic Acids-DNA Primary Structure

The primary structure of a nucleic acid is its covalent structure and nucleotide sequence.

The back bone of the primary structure is the linear strand made by sugar phosphate residues, linked together, while the bases project laterally. This way a long, un-branched chain is formed.

Primary structure is a huge linear polymer of dNTPs that are joined to



The resulting long, un-branched chain has polarity.

At one end there is a free 5' carbon (phosphate group) and at the other end there is a free 3' carbon (hydroxyl group) that are not attached to other nucleotides.

The variable part is concerned with the expression of genetic information.

"

By convention, the structure of a single strand of nucleic acid is always written with the 5' end on the left and the 3' end on the right. For example, the sequence 5'-ATCG-3' is written as ATCG.

3' "Zywo"•swzvo°oz°o•ox'k'syx•"yp'r 9 "zox'k"noy—"s yx#moy'sno"ko° " "zH-C-G-T-H"VO1" " pApCpGpTpA, and finally 5'-ACGTA-3'.

174 Nucleotides and Nucleic Acids-DNA Secondary Structure

DNA Secondary Structure

Any regular, stable structure taken up by some or all of the nucleotides in a nucleic acid can be referred to as secondary structure

DNA exist as double stranded molecule (double helix), except few viruses that contain single stranded molecule.

This model was presented by Watson and Crick in 1953.

The two polydeoxyribonucleotide strands are coiled around a common axis called axis of symmetry.

The chains are paired in anti-parallel fashion. The 5'-end of one strand is paired with the 3'-end of the other strand.

In the DNA helix

Deoxyribose-phosphate backbone is hydrophilic and thus it is on the outside of the molecule.

The bases are hydrophobic and are thus stacked inside the molecule.

The overall structure resembles a twisted ladder.

Grooves: The spatial relationship between the two strands in the helix creates
a major (wide) groove
a minor (narrow) groove.

These grooves provide access for the binding of regulatory proteins to their specific recognition sequences along the DNA

175 Nucleotides and Nucleic Acids-DNA Tertiary Structure

DNA tertiary Structure

The complex folding of large chromosomes within eukaryotic chromatin and bacterial nucleoids is generally considered tertiary structure

Eukaryotic DNA is associated with tightly bound basic proteins, called histones.

These serve to order the DNA into fundamental structural units, called nucleosomes.

There are five classes of histones, designated H1, H2A, H2B, H3, and H4.

These are positively charged at physiologic pH as a result of their high content of lysine and arginine.

Two molecules each of H2A, H2B, H3, and H4 form the structural core of the nucleosome.

Around this core, a segment of the DNA double helix is wound nearly twice approximately 140bp

The DNA wrapped around the nucleosome core is continuous and joins one nucleosome core to the next -the linker DNA

this 50 bp DNA is complexed with the fifth type of histone, H1.

Nucleosomes can be packed more tightly to form a polynucleosome also called a nucleofilament or a 30-nm fiber.

The fiber is organized into loops that are anchored by nuclear scaffolding proteins.

Additional levels of organization lead to the final chromosomal structure

176 Nucleotides and Nucleic Acids-DNA Properties

DNA Properties

Template and Non-template Strands

The term template strand refers to the sequence of DNA that is copied during the synthesis of mRNA.

The opposite strand is called the Non Template or coding strand or the mRNA-like strand

it has base sequence directly corresponding to the mRNA sequence

the sequence corresponds to the codons that are translated into protein

The 3'-5' strand is called Template strand while

5'-3' strand is called Non Template (coding strand).

Base Pairing

The bases of one strand of DNA are paired with the bases of the opposite strand, so that

Adenine is always paired with thymine

Cytosine is always paired with guanine.

One polynucleotide chain of DNA double helix is always complementary to the other

Given the sequence of bases on one chain, the sequence of bases on the complementary chain can be determined

177 Nucleotides and Nucleic Acids-DNA Properties Contd

J r k°qkpp•"Y≠w•"

Due to specific base pairing of DNA i.e A toT and G to C

In any sample of double-stranded DNA the amount of adenine equals the amount of thymine.

The amount of guanine equals the amount of cytosine

the total amount of purines equals the total amount of pyrimidines ;

i.e. $A + G = T + C$.

The base pairs are held together by hydrogen bonds

Two between A and T

Three between G and C

These hydrogen bonds, plus the van der Waals and hydrophobic interactions between the adjacent stacked bases, stabilize the structure of the double helix. The base composition of DNA varies from one species to another, but is same if isolated from different tissues of same species. The base composition does not change with age, nutritional status and environment.

178 Denaturation of DNA

DNA Denaturation

Separation of the two strands of the double helix when hydrogen bonds between the paired bases are disrupted.

Disruption can occur in the laboratory if

the pH or the salt concentration of the DNA solution is altered

if the solution is heated above 80°C

Disruption of the hydrogen bonds between paired bases and of base stacking causes unwinding of the double helix to form two single strands complete separation of DNA stands along the entire length.

No covalent bonds in the DNA are broken

RNA duplexes are more stable than DNA duplexes.

At neutral pH denaturation of a double helical RNA often requires temperatures 20° C or more higher than those required for denaturation of a DNA molecule with a comparable sequence.

T_m: When DNA is heated, the temperature at which one half of the helical structure is lost is defined as the melting temperature.

A specific sequence of DNA has a characteristic denaturation temperature, or melting point (T_m)

Careful determination of the melting temperature of a DNA specimen, under fixed conditions of pH and ionic strength, can yield an estimate of a DNA base composition.

179 Nucleotides and Nucleic Acids-II

DNA Denaturation (Contd.)

Concomitant with this denaturation of the DNA molecule is an increase in the optical absorbance of the purine and pyrimidine bases a phenomenon referred to as hyperchromicity of denaturation.

So single-stranded DNA has a higher relative absorbance at this wavelength than does double-stranded DNA.

Denaturation can be monitored by measuring its absorbance at 260 nm

Factors affecting T_m

The T_m is influenced by

the base composition of the DNA

the salt concentration of the solution

The higher the content of GC base pairs, the higher the melting point of the DNA.

This is because GC base pairs, with three hydrogen bonds, require more heat energy to dissociate than AT base pairs.

An *increase* in salt concentration *increases* and a decrease in salt concentration decreases the T_m

180 Nucleotides and Nucleic Acids-DNA Renaturation

DNA Renaturation

Under appropriate conditions (*temp. & salt concentration*), separated strands of DNA will renature or reassociate and form the double helix by the process called renaturation (or reannealing).

This reannealing process is also referred to as hybridization.

When the temperature or pH is returned to the range in which most organisms live, the unwound segments of the two strands spontaneously rewind, or anneal, to yield the intact duplex

Renaturation of a DNA molecule is a rapid one-step process.

The rate of reassociation depends upon the concentration of the complementary strands.

At a given temperature and salt concentration, a particular nucleic acid strand will associate tightly only with a complementary strand

This property is utilized in analyzing nucleotide sequencing of a given nucleic acid.

Under appropriate conditions DNA will form a hybrid with a complementary DNA or with a complementary RNA

Hybridization is combined with gel electrophoresis techniques that separate nucleic acids by size,

coupled with radioactive or fluorescent probe labeling to provide a detection of a nucleotide sequence.

181 Nucleotides and Nucleic Acids- RNA

RNA

The genetic master plan is contained in the nucleotide sequence of DNA.

P's "r °y≠qr "r o"°\$ yx≠mosrkns"-YUH. 'r o"&Δy°usq"nyzs•&yp"r o"-KUH." "rk""r o" master plan is expressed

RNA is a polymer of ribonucleotides of Adenine, Uracil, Guanine and Cytosine, joined together by 3'-5' phosphodiester bonds.

RNA does not contain thymine except in rare cases.

The pentose sugar of RNA is D-ribose.

Location: RNA is found in the nucleolus, ribosomes, mitochondria, and cytoplasm.

The genetic material for some animal and plant viruses is RNA rather than DNA.

There is a wide variety of RNAs,

messenger RNAs (mRNAs)- transfer genetic information from DNA to the proteinsynthesizing machinery.

ribosomal RNAs (rRNAs)- contribute to the formation and function of ribosomes

transfer RNAs (tRNAs)- adapter molecules that carry specific amino acids for protein synthesis

small nuclear RNA (snRNA)- play pivotal roles in RNA processing, particularly mRNA processing

Due to their intrinsic catalytic activity these RNA enzymes, are called ribozyme

182 Nucleotides and Nucleic Acids-Difference between RNA and DNA

Differences between DNA and RNA

Although sharing many features with DNA, RNA possesses several specific differences.

Size: They are considerably smaller than DNA

Sugar: In RNA, the sugar moiety to which the phosphates and purine and pyrimidine bases are attached is ribose rather than the 2'-deoxyribose. Pyrimidine: The pyrimidine components of RNA differ from those of DNA.

Instead of thymine, RNA contains the ribonucleotide of uracil.

Thymine is present in the rare case of tRNA

Single Strand:

RNA typically exists as a single strand whereas DNA exists as a double-stranded helical molecule.

However, given the proper complementary base sequence with opposite polarity, the single strand of RNA is capable of self- folding like a hairpin and thus acquiring doublestranded characteristics: G pairing with C, and A with U.

Chargaff's Rules Do Not Apply:

Since the RNA molecule is a single strand complementary to only one of the two strands of a gene,

N J "

H \ "

~~W^oxo"~~ "W^oswmxo"

Hydrolysis: RNA can be hydrolyzed by alkali to 2',3' cyclic diesters of the mononucleotides, compounds that cannot be formed from alkali-treated DNA because of the absence of a 2'-hydroxyl group.

Location: In addition to nucleus, RNA is found in cytoplasm.

Reverse Transcription: DNA forms RNA by transcription whereas the process by which RNA form DNA is called reverse transcription.

183 Nucleotides and Nucleic Acids- RNA Structural Hierarchy

RNA Structural Hierarchy

RNA has no simple, structural hierarchy that serves as a reference point, as does the double helix for DNA.

The three-dimensional structures of many RNAs, like those of proteins, are complex and unique.

Primary Structure of RNA

It is defined as the number and sequence of ribonucleotides in the RNA chain.

The sequence is complementary to the template strand of the gene from which it was transcribed.

template strand, the sequence of DNA that is copied.

Non Template (coding strand).

The ribonucleotides are held together by 3' - 5' phosphodiester bonds.

3'-OH group of one nucleotide is bound to 5'-PO₄ of the other nucleotide and form a linear strand.

The ribosyl moieties are attached to the nucleobases by *N*-glycosidic bonds "

Similar to DNA RNA polymer also has polarity

184 Nucleotides and Nucleic Acids- Secondary Structure of RNA

Secondary Structure of RNA

Secondary structure involves coil formation of the polyribonucleotide chain.

The coiled structures are stabilized by Hydrophobic interactions between purine and pyrimidine bases. Intra-chain hydrogen bonds between G-C and A-U.

Note: Adenine pairs with Uracil in RNA 

Weak interactions, especially base-stacking interactions.

Where complementary sequences are present, the predominant double-stranded structure is formed.

Common RNA secondary structures

Internal loops: a short series of unpaired bases in a longer paired helix and

bulges: regions in which one strand of a helix has "extra" inserted bases with no counterparts in the opposite strand

Tertiary Structure of RNA

It is the folding of the molecule into three dimensional structure.

cross-linking at various sites stabilized by hydrophobic and hydrogen bonds produces a compactly coiled globular structure.

The stacking of helices, together with specific helix helix contacts or helix loop interactions, lead to compact tertiary structure of the RNA assemblies, generally in the presence of divalent ions or polyamines

185 Nucleotides and Nucleic Acids- Messenger RNA

Messenger RNA (mRNA)

This class is the most heterogeneous in

Abundance

Size (500-6000 nucleotides)

base sequence

Stability

mRNA comprise about 2-5% of total cellular RNA

mRNA molecules are formed with the help of DNA template strand (3'-5') during the process called transcription.

In addition to the protein coding regions in the mature eukaryotic mRNA that can be translated,

there are untranslated regions at its 5' and 3' ends

Moreover, there is a 5' cap and

a poly A tail at 3' end

Function of mRNA

The members of this class function as messengers to convey the information in a gene to the protein synthesizing machinery.

The mRNA carries genetic information from the nuclear DNA to the cytosol, where it is used as a template for protein synthesis.

186 Nucleotides and Nucleic Acids- Transfer RNA

Transfer RNA (tRNA)

tRNA is the smallest of the three major species of RNA (4S).

They are single stranded globular molecules.

They remain largely in cytoplasm.

They are generated by nuclear processing of a precursor molecule.

tRNAs compose roughly 20% of total cellular RNA

There are at least 20 species of tRNA molecules in every cell.

Although each specific tRNA differs from the others in its sequence of nucleotides, the tRNA molecules as a class have many features in common

Primary structure

t RNA molecules consist of 74-95 nucleotides in a particular sequence.

The t RNA molecules contain not only the usual bases like adenine, guanine, cytosine, uracil but also contain **unusual bases**

These unusual bases(also called modified bases) include

Ds -n°y°kns" " Pseudouridine

Thymine.

Secondary Structure Pseudouridine

Each single stranded

t RNA is folded extensively.

Extensive intra chain base pairing which leads to a characteristic CLOVER-LEAF structure.

These folds are stabilized by hydrogen bonds between complementary bases of the same strand.

187 Nucleotides and Nucleic Acids- Transfer RNA (continued..)

Arms or loops of tRNA

All tRNA molecules contain 4 main arms or loops.

1-Acceptor arm: This is made up of unpaired sequences of cytosine-cytosine-adenine (CCA) at the 3' end.

The 3'OH group of adenine binds with the carboxylic group of a specific amino acid and carries it to ribosomes for protein synthesis.

2-Anticodon arm: It is in the form of a loop and carries specific sequences of three bases which constitute the anticodon.

The bases of anticodon are bonded with three complementary bases of codon on mRNA.

3-D arm: It contains the base dihydrouridine.

4-Z I 'j °v ? It contains thymine, pseudouridine and cytosine.

[r o"o—°k"kw"kn"r o" J "kw•"r ov"y"nopxok"•zonsp"YUH3'

D arm: contains dihydrouridine. **Z I 'j °v ?** thymine, pseudouridine and cytosine.

Function of tRNA

The t RNA molecules serve as ADAPTERS for the translation of information in the sequence of nucleotides of the mRNA into specific amino acids.

There is at least one (and often several) specific type of tRNA molecule for each of the amino acids commonly found in proteins.

Each t RNA carries its specific amino acid to the site of protein synthesis.

There it recognizes the genetic code word on mRNA (codon) and this specifies the addition of its amino acids to the growing peptide chain.

188 Nucleotides and Nucleic Acids- TRNA Continued

189 Nucleotides and Nucleic Acids- ribosomal RNA

Ribosomal RNA (rRNA)

found in association with several proteins as a component of ribosomes--- a cytoplasmic nucleoprotein structure that acts as the machinery for the synthesis of proteins from the mRNA template.

RNAs make up 80% of the total RNA in the cell.

The ribosomal subunits are defined according to their sedimentation velocity in Svedberg units.

Svedberg unit is related to the molecular weight and shape of the compound.

The bases in r RNA are mainly adenine, guanine, cytosine and uracil and a few pseudouridine

Eukaryotic Ribosome

The mammalian ribosome contains two major nucleoprotein subunits:

- a larger one with 60S

- a smaller one with 40S.

The **60S** subunit contains

- a 5S rRNA

- a 5.8S rRNA

- a 28S rRNA

- more than 50 specific polypeptides.

The **40S** subunit is smaller and contains

- a single 18S rRNA

- Approx. 30 distinct polypeptide chains.

'''

'''

In eukaryotes, all of the ribosomal RNA molecules except the 5S rRNA, which is independently transcribed, are processed from a single 45S precursor RNA molecule in the nucleolus- packed with the specific ribosomal proteins.

The rRNA are necessary for ribosomal assembly and play a key role in the binding of

mRNA to ribosomes and its translation

In the cytoplasm, the ribosomes remain quite stable and capable of many translation cycles

190 Nucleotides and Nucleic Acids- small Nuclear RNA

Other types of RNA

Small nuclear RNA (snRNA)

Small nuclear RNA (snRNA) are large number of small stable RNA species found in eukaryotic cells.

Most of them are complexed with proteins to form ribonucleoproteins.

They are distributed in the nucleus, in the cytoplasm or in both.

They are significantly involved in rRNA and mRNA processing and gene regulation.

Large & Small Noncoding Regulatory RNAs

One of the most exciting discoveries in the last decade of eukaryotic regulatory biology has been the identification and characterization of regulatory nonprotein coding RNAs (ncRNAs).

NcRNAs exist in two general size classes,

small consisting of microRNA (miRNAs) and silencing (siRNAs) and

Large consisting of long noncoding RNAs (lncRNAs)

The small ncRNAs termed microRNA (miRNAs) and silencing (siRNAs) typically inhibit gene expression at the level of specific protein production by

targeting mRNAs through one of several distinct mechanisms.

Both siRNAs and miRNAs typically hybridize, via the formation of RNA-RNA hybridization to their targeted mRNAs

long noncoding RNAs (lncRNAs).

lncRNAs, which as their name implies, do not code for protein (ie, the mRNA encoding genes).

ncRNAs make up a significant portion of eukaryotic transcription

ncRNAs play many roles ranging from contributing to structural aspects of chromatin to regulation of mRNA gene transcription by RNA polymerase II.

191 Enzymes

Reaction catalysts of biological systems: the enzymes, the most remarkable and highly specialized proteins.

Enzymes have extraordinary catalytic power, often far greater than that of synthetic or inorganic catalysts.

With the exception of some catalytic RNA molecules all enzymes are proteins.

Thus the primary, secondary, tertiary, and quaternary structures of protein enzymes are essential to their catalytic activity. ☰

Enzyme-catalyzed reactions have three basic steps:

binding of substrate: $E + S \rightarrow ES$

conversion of bound substrate to bound product: $ES \rightarrow EP$

release of product : $EP \rightarrow E + P$

They have a high degree of specificity for their substrates, they function in aqueous solutions under very mild conditions of temperature and pH.

Enzymes do not invent new reactions; they simply make reactions occur faster.

In addition to increasing the speed of reactions, enzymes provide means for regulating the rate of metabolic pathways in the body.

The commonly used names for most enzymes describe the type of reaction catalyzed, followed by the suffix **ase** e.g.

dehydrogenases remove hydrogen atoms

proteases hydrolyze proteins

Modifiers may precede the name to indicate, for example,
the substrate (xanthine oxidase),
the source of the enzyme (pancreatic ribonuclease),
its regulation (hormone-sensitive lipase)

Where needed, alphanumeric designators are added to identify multiple forms of an enzyme e.g,

RNA polymerase III

protein kinase C

192 Enzymes- classification

IUB Classification of Enzymes

International Union of Biochemists (IUB) developed an unambiguous system of enzyme nomenclature in which each enzyme has a

unique name and

code number

As an example, the formal systematic name of the enzyme (hexokinase) catalyzing the reaction

$\text{ATP} + \text{D-glucose} \rightarrow \text{ADP} + \text{D-glucose-6-phosphate}$ is

ATP:glucose phosphotransferase,

Its Enzyme Commission (E.C.) number is 2.7.1.1.

(2) denotes the class name (transferase)

(7) the subclass phosphotransferase

(1) denotes a hydroxyl group as acceptor

(1) D-glucose as the phosphoryl group acceptor.

In the systematic naming system, enzymes are divided into six major classes each with numerous subgroups

1) Oxidoreductases

catalyze oxidation reduction reactions

further divided into four subgroups;

Oxidase,

Hydroperoxidases

Oxygenases.

2) Transferases

These bring about a transfer of functional groups such as phosphate and amino group from one molecule to another molecule called donor and acceptor molecules respectively.

The common examples of this group are

Transminases

Phosphotransfrases (Kinases)

Hexokinase is a phosphotransfease which catalyze the transfer of phosphate groups.

$\text{NAD}^+ + \text{glucose} + \text{P}_i \rightarrow \text{NADH} + \text{glucose-6-phosphate} + \text{ADP}$.

193 Enzymes- IUB classification (continued..)

3) Hydrolases

These enzymes catalyze hydrolysis, i.e.

add water molecule to the substrate which is simultaneously decomposed; the functional group of substrate is transferred to water.

Common example of hydrolases are:

Protein hydrolyzing Enzymes (peptidases).

Carbohydases

Lipid hydrolyzing enzymes e.g. Lipases and

Phospholipases.

4) Lyases

These enzymes catalyze the addition of

NH₂,

H₂O or

CO₂ to double bonds or

the removal of these groups leaving behind double bonds.

Lyases are included in a separate class because they catalyze these reactions by means other than hydrolysis or oxidation.

194 Enzymes- IUB Classification (Continued...)

5) Isomerases

These enzymes catalyze the structural change within a single molecule by the transfer of groups within it, resulting in the formation of an isomeric form of substrate.

6) Ligases

These enzymes catalyze condensation reactions joining two molecules by forming

C-O,

C-S,

C-N and

C-C bonds.

The energy for condensation is provided by cleavage of high energy phosphates, e.g. ATP, GTP etc.

195 Enzymes- Ligand

Ligand

In biochemistry and pharmacology, a **ligand** (from the Latin *ligandum*, *binding*) is a substance (usually a small molecule),

that forms a complex with a biomolecule to serve a biological purpose.

In a narrower sense, it is a signal triggering molecule, binding to a site on a target protein

A molecule bound reversibly by a protein is called a **ligand**.

Ligands include substrates, inhibitors, activators, and neurotransmitters

A ligand may be any kind of molecule, including another protein.

A ligand binds at a site on the protein called the **binding site**, **binding site** is complementary to the ligand in size, shape, charge, and hydrophobic or hydrophilic character.

Furthermore, the interaction is **specific**: the protein can discriminate among the thousands of different molecules in its environment and selectively bind only one or a few.

The binding of a protein and ligand is often coupled to a **conformational change** in the protein that makes the binding site more complementary to the ligand, permitting tighter binding called **induced fit**.

A given protein may have **separate binding sites** for several different ligands.

In a multi-subunit protein, a conformational change in one subunit often affects the conformation of other subunits.

Interactions between **ligands** and proteins may be regulated, through interactions with additional ligands.

These other ligands may cause conformational changes in the protein that affect the binding of the **first ligand**.

196 Enzymes- Mechanism of enzyme action

Mechanism of Enzyme Action

Enzymes bind and chemically transform other molecules they catalyze reactions.

The molecules acted upon by enzymes are called reaction substrates rather than ligands.

"

Enzymes are highly effective catalysts, commonly enhancing reaction rates by a factor of 10^5 to 10^{17} .


The distinguishing feature of an enzyme-catalyzed reaction is that it takes place within the confines of a pocket on the enzyme called the **active site**.

The surface of the active site is lined with **amino acid** residues with side chains that complement and bind the substrate and catalyze its chemical transformation.

Often, the active site encloses a substrate, sequestering it completely from solution.

The enzyme substrate complex, whose existence is central to the action of enzymes.

Enzyme-catalyzed reactions have three basic steps:




To understand catalysis, we must first appreciate the important distinction between reaction equilibria and reaction rates.

197 Enzymes- Mechanism of enzyme action (continued..)

Reaction Equilibria

The function of a catalyst is to increase the rate of a reaction.

Catalysts *do not affect reaction equilibria*



"

A diagram of this kind is a description of the energy changes during the reaction.

The horizontal axis (reaction coordinate) reflects the progressive chemical changes (e.g., bond breakage or formation) as S is converted to P.

Reaction coordinate diagram for a chemical reaction.

The free energy of the system is plotted against the progress of the reaction $S \rightarrow P$. A diagram of this kind is a description of the energy changes during the reaction, and the horizontal axis (reaction coordinate) reflects the progressive chemical changes (e.g., bond breakage or formation) as S is converted to P. The activation energies, $G^\ddagger$, for $S \rightarrow P$ and $P \rightarrow S$ reactions are indicated. G° is the overall standard free-energy change in the direction $S \rightarrow P$.

Activation energies, $G^\ddagger$

The activation energies, $G^\ddagger$, for the

$S \rightarrow P$ and $P \rightarrow S$ reactions are indicated.

G° is the overall standard free-energy change in the direction $S \rightarrow P$.

Free-energy change, G°

When a reacting system is not at equilibrium, the tendency to move toward equilibrium represents a driving force

the magnitude of which can be expressed as the free-energy change for the reaction, G° .

Standard free-energy change, G°

Under standard conditions (298 K = 25 °C)

when reactants and products are initially present at 1 M concentrations or

for gases, at partial pressures of 101.3 (kPa), or 1 atm

the force driving the system toward equilibrium is defined as the standard free-energy change, G° .

However, because the conditions in the body systems are different from standard conditions

energy in biological systems is described in terms of free energy, G° .

198 Enzymes-Mechanism of action (continued...)

Ground State & Transition State

The starting point for either the forward or the reverse reaction is called the ground state

The equilibrium between S and P reflects the difference in the free energies of their ground states.

The free energy of the ground state of P is **lower** than that of S

So G° for the reaction is negative and the equilibrium favors P.

The position and *direction* of equilibrium are *not affected by* any catalyst.

Reaction coordinate diagram for a chemical reaction.

The free energy of the system is plotted against the progress of the reaction

The vertical axis (free energy) reflects the progressive chemical changes (e.g., bond breakage or formation) as S is converted to P. The activation energies, $G^\ddagger$, are the energy difference between the ground state and the transition state. G° is the overall standard free-energy change for the reaction. The energy barrier between S and P: The energy required for alignment of reacting groups, formation of transient unstable charges, bond rearrangements and other transformations

[r s "s "sw• °k'on"l —"r o"oxo°q—"r sw"

To undergo reaction, the molecules must overcome this barrier and therefore must be raised to a higher energy level

This is called the **transition state**.

It is simply a fleeting molecular moment in which events such as bond breakage, bond formation, and charge development have proceeded to the precise point at which decay to either substrate or product is equally likely.

The difference between the energy levels of the ground state and the transition state is the **activation energy, $G^\ddagger$**

At the top of the energy hill is a point at which decay to the S or P state is equally probable (it is downhill either way)

A substance that modifies the transition state to lower the activation energy is termed a catalyst; a biological catalyst is termed an enzyme.

199 Enzymes-Mechanism of Action- Activation Energy (continued...)

Mechanism of Enzyme Action (Contd.)

Hm's/k'syX"oxo°q—" $G^\ddagger$.

The difference between the energy levels of the ground state and the transition state is $G^\ddagger$.

The rate of a reaction reflects this activation energy: a higher activation energy corresponds to a slower reaction.

Reaction rates can be increased by raising the temperature, thereby increasing the number of molecules with sufficient energy to overcome the energy barrier.

Alternatively, the activation energy can be lowered by adding a catalyst.

Catalysts enhance reaction rates by lowering activation energies.

Reaction coordinate diagram comparing enzyme catalyzed

and uncatalyzed reactions. E_a^{uncat} and E_a^{cat}

and EP intermediates occupy minima in the energy progress curve of

the enzyme-catalyzed reaction. The terms G^{uncat}

uncat and G^{cat} are the activation energies for the uncatalyzed

reaction and the overall activation energy for the catalyzed reaction, respectively.

The activation energy is lower when the enzyme catalyzes the reaction.

The role of enzymes is to accelerate the inter-conversion of S and P.

i.e. **enzymes lower the energy of activation, G^{cat} of a reaction.**

The enzyme is not used up in the process, and the equilibrium point is unaffected.

However, the reaction reaches equilibrium much faster when the appropriate enzyme is present,

because the rate of the reaction is increased.

200 Enzymes- induced fit hypothesis

The Induced Fit Hypothesis

Some proteins can change their shape (conformation)

to form a shape that is complementary to the substrate. This change in conformation is induced by multiple weak interactions with the substrate.

This change in conformation when the substrate binds is induced by multiple weak interactions with the substrate.

There may also be rearrangements of covalent bonds during an enzyme-catalyzed reaction.

This conformational change is referred to as induced fit.

The Induced Fit Hypothesis

Chemical reactions of many types take place between substrates and enzyme • "functional groups (specific amino acid side chains, metal ions, and coenzymes

Induced fit serves to bring specific functional groups on the enzyme into the proper position to catalyze the reaction.

Induced Fit Theory

Enzyme is not rigid, changes shape with substrate.

The active site is also moulded into a precise conformation

Making the chemical environment suitable for the reaction

201 Enzymes- co factors, coenzymes

Cofactors, Coenzymes and Prosthetic groups

Some enzymes require no chemical groups for activity other than their amino acid residues.

Whereas some enzymes require molecules other than proteins for enzymic activity.

If the non-protein moiety is a metal ion such as Zn^{2+} or Fe^{2+} , it is called a cofactor.

If it is a complex organic molecule or metallo-organic compound it is termed a coenzyme.

A coenzyme or metal ion that is very tightly or even covalently bound to the enzyme protein is called a prosthetic group.

The term holoenzyme refers to the active enzyme with its non-protein component,

whereas the enzyme without its non-protein moiety is termed an apoenzyme (apoprotein) and is inactive.

Coenzymes serve as recyclable shuttles that transport many substrates from one point within the cell to another.

The function of these shuttles is twofold.

First, they stabilize species

such as hydrogen atoms (FADH) or hydride ions (NADH)

that are too reactive to persist for any significant time in the presence of the water or organic molecules that permeate cells.

Second, they serve as an adaptor or handle that facilitates the

recognition and binding of small chemical groups, such as acetate (coenzyme A) or glucose (UDP), by their target enzymes

202 Enzymes-Reaction rates and order

Reaction Velocity (v)

The rate or velocity of a reaction (v) is the number of substrate molecules converted to product per unit time;

Velocity is usually expressed as **μmol of product** formed **per minute**.

The rate of an enzyme-catalyzed reaction increases with substrate concentration until a maximal velocity (V_{max}) is reached- reflecting the **saturation** with substrate of **all available binding sites are occupied** on the enzyme molecules present.

(V_{max}) is reached- reflecting the **saturation** with substrate of **all available binding sites are occupied** on the enzyme molecules present.

The **equilibrium constant, K_{eq}**, or simply **K**.

Under the standard conditions used to compare biochemical processes, an equilibrium constant is denoted **K_{eq}** (or **K**).

The rate of any reaction is determined by the concentration of the reactant (or reactants) and by a **rate constant**, usually denoted by **k**.

The amount of S that reacts per unit time is expressed by a **rate equation**: $V = k[S]$.

In this reaction, the rate depends only on the concentration of S.

This is called a first-order reaction.

The factor k is a proportionality constant that reflects the probability of reaction under a given set of conditions (pH, temperature, and so forth).

If a reaction rate depends on the concentration of two different compounds.

or if the reaction is between two molecules of the same compound, the reaction is second order.

The rate equation then becomes

$$V = k[S][S]$$

203 Enzyme-Factor affecting enzyme activity

Enzymes can be isolated from cells, and their properties studied in a test tube (that is, in vitro).

Factors Affecting Enzymatic Activity

Different enzymes show different responses to changes in;

substrate concentration

temperature, and

pH.

Enzymic responses to these factors give us valuable clues as to how enzymes function in living cells (that is, in vivo)

Substrate concentration

The rate of an enzyme-catalyzed reaction increases with substrate concentration until maximal velocity (V_{max}) is reached

The leveling off of the reaction rate at high substrate concentrations reflects the saturation with substrate of all available binding sites on the enzyme molecules present.

Temperature

The reaction velocity increases with temperature until a peak velocity is reached

This increase is the result of the increased number of molecules having sufficient energy to pass over the energy barrier and form the products.

Further elevation of the temperature results in a decrease in reaction velocity as a result of temperature-induced denaturation of the enzyme

204 Enzymes-Factors (continued..)

Effect of pH

The pH optimum varies for different enzymes:

The pH at which maximal enzyme activity is achieved is different for different enzymes, and often reflects the $[H^+]$ at which the enzyme functions in the body.

For example, pepsin, a digestive enzyme in the stomach, is maximally active at pH 2.

Whereas other enzymes, designed to work at neutral pH, are denatured by such an acidic environment

Another examples is that there are two types of phosphatases in the body.

The one that acts in the alkaline pH is called alkaline phosphatase and

the other which acts at acidic pH is known as acid phosphatase.

Effect of pH on the ionization of the active site:

The concentration of H^+ affects reaction velocity in several ways.

First, the catalytic process usually requires that the enzyme and substrate have specific chemical groups in either an ionized or un-ionized state in order to interact.

For example, catalytic activity may require that an amino group of the enzyme be in the protonated form (NH_3^+).

At alkaline pH, this group is deprotonated, and the rate of the reaction, therefore, declines.

Extremes of pH can also lead to denaturation of the enzyme.

205 Enzymes-Enzyme kinetics

Enzyme Kinetics

Substrate Concentration Affects the Rate of Enzyme-Catalyzed Reactions

L "O"Z" "LZ" "LW"

"L "O"W"

A key factor affecting the rate of a reaction catalyzed by an enzyme is the concentration of substrate, $[S]$.

However, studying the effects of substrate concentration is complicated because substrate is converted to product

and because of reversibility of reactions, e.g. conversion of product back to substrate.

One simplifying approach in kinetics experiments is to measure the **initial rate (or initial velocity), designated V_0** , when $[S]$ is much greater than the concentration of enzyme,

$[E]$.

The velocity (v) of a reaction is the rate of product formation

Whereas V_0 is the initial velocity and is measured as soon as the reactants and enzymes are mixed.

At the start of a reaction, $[S]$ is in large excess of $[P]$.

Thus the initial velocity of the reaction will be dependent on substrate concentration

At that time, the concentration of product is very small and, therefore, the rate of the back reaction from P to S can be ignored.

Increase in substrate concentration increases V_0 i.e.

initial velocity is increased whenever a fixed concentration of enzyme is mixed with an increased concentration of substrate.

206 Enzymes-Enzyme kinetics (continued..)

Maximal Velocity V_{max}

When initial velocity (V_0) is plotted against $[S]$,

a **hyperbolic curve** results,

where V_{max} represents the maximum reaction velocity.

At relatively low concentrations of substrate, V increases almost linearly.

At higher substrate concentrations, V increases by smaller and smaller amounts in response to increases in $[S]$.

This plateau-like V region is close to the **maximal velocity, V_{max}** .

V_{max} is extrapolated from the plot, because V approaches but never quite reaches V_{max} .

At this point in the reaction, if $[S] \gg E$, all available enzyme is "saturated" with bound substrate, meaning only the **ES complex** is present.

This dictates that further increasing the substrate concentration will not result in increased V_0

A. Low $[S]$ B. 50% $[S]$ or K_m C. High, saturating $[S]$

The leveling off of the reaction rate at high substrate concentrations reflects the saturation with substrate of all available binding sites on the enzyme molecules present.

This finite limit of V_{max} is called saturation kinetics.

Saturation kinetics is a characteristic property of all rate processes dependent on the binding of a ligand to a protein e.g. mThe process by which a **carrier protein** transfers a **solute molecule** across the **lipid bilayer** resembles an **enzyme-substrate reaction**, and in many ways carriers behave like enzymes. In contrast to ordinary enzyme-substrate reactions, however, the transported solute is not covalently modified by the carrier protein, but instead is delivered unchanged to the other side of the **membrane**.

embrane transporter proteins.

207 Enzymes-Enzyme kinetics K_m (continued...)

Michaelis constant K_m

The substrate concentration at which V is half maximal is K_m , the Michaelis constant.

The effect of varying $[S]$, on V_0 , when the enzyme concentration is held constant

K_m reflects the affinity of the enzyme for the substrate.

Small K_m reflects a high affinity of the enzyme for substrate, because a low concentration of substrate is needed to half-saturate the enzyme i.e. to reach a velocity $\frac{1}{2} V_{max}$. A numerically large (high) K_m reflects a low affinity of enzyme for substrate because

a high concentration of substrate is needed to half-saturate the enzyme.

- a. **Small K_m :** A numerically small (low) K_m reflects a high affinity of the enzyme for substrate, because a low concentration of substrate is needed to half-saturate the enzyme $\frac{1}{2} V_{max}$.
- b. **Large K_m :** A numerically large (high) K_m reflects a low affinity of enzyme for substrate because a high concentration of substrate is needed to half-saturate the enzyme
- c. The velocity of an enzyme is most sensitive to changes in substrate concentration over a concentration range below its K_m .
- d. At substrate concentrations less than K_m , doubling the substrate concentration nearly doubles the velocity of the reaction
- e. At substrate concentrations 10 times the K_m , doubling the substrate concentration has little effect on the velocity.

A comparison between the isozymes of hexokinase illustrates the significance of the K_m .

Hexokinase catalyses the first step in glucose metabolism in most cells, the transfer of a phosphate from ATP to glucose to form glucose 6-phosphate.

Hexokinase I, the isozyme in red blood cells has a low K_m for glucose of approximately 0.05 mM- helpful in utilizing blood glucose even when the blood glucose concentration is very low.


The isozyme of hexokinase, called glucokinase, which is found in the liver has a much higher K_m of approximately 5 to 6 mM- responsible for storing glucose as glycogen or converting it to fat after a carbohydrate meal.

208 Enzymes-Michaelis menten equation

Michaelis-Menten equation, the rate equation

Leonor Michaelis and Maud Menten in 1913, proposed a simple model that accounts for most of the features of enzyme-catalyzed reactions.

They postulated that the enzyme first combines reversibly with its substrate to form an enzyme-substrate complex in a relatively fast reversible step:



$K-1$

The ES complex then breaks down in a slower second step to yield the free enzyme (E) and the reaction product (P):

$K-2$



$K-2$


Early in the reaction, the concentration of the product, **[P], is negligible**, and we make the assumption that the reverse reaction ($K-2$), can be ignored

This assumption is not critical but it simplifies our task.

The overall reaction then reduces to



$K-1$



E = Enzyme S = Substrate P = Product

ES = Enzyme-Substrate complex k_1 rate

constant for the forward reaction

k_{-1} = rate constant for the breakdown of the ES to substrate k_2

= rate constant for the formation of the products

Because the slower second reaction must limit the rate of the overall reaction.

The overall rate must be proportional to the concentration of the species that react in the second step, that is, ES.

The maximum initial rate of the catalyzed reaction (V_{max}) is observed when virtually all the enzyme is present as the ES complex and [E] is vanishingly small.

Under these conditions, increases in $[S]$ have no effect on rate.

This condition exists when $[S]$ is sufficiently high that essentially all the free enzyme has been converted to the ES form.

After the ES complex breaks down to yield the product P, the enzyme is free to catalyze reaction of another molecule of substrate.

209 Enzymes-Michaelis Menten equation (continued..)

Michaelis-Menten equation, the rate equation (Contd.)

$$V = \frac{V_{\max} [S]}{K_m + [S]}$$

$$K_m + [S]$$

Where;

V = initial reaction velocity

$V_{\max}$ = maximal velocity

K_m = Michaelis constant

$$= \frac{K_1 + K_2}{K_{-1}}$$

$[S]$ = substrate concentration

This is the **Michaelis-Menten equation**-statement of the quantitative relationship between the;

initial velocity V ,

the maximum velocity $V_{\max}$, and

the initial substrate concentration $[S]$, all related through the Michaelis constant K_m

An important numerical relationship emerges from the Michaelis-Menten equation in the special case;

when V is exactly one-half $V_{\max}$

$$V = \frac{V_{\max} [S]}{K_m + [S]}$$

$$K_m + [S]$$

$$V_{\max} = \frac{V_{\max} [S]}{K_m + [S]}$$

$$2 = \frac{K_m + [S]}{[S]}$$

On dividing by $V_{\max}$, we obtain

$$1/2 = [S] / K_m + [S]$$

Solving for K_m , we get

$$K_m + [S] = 2[S], \text{ or}$$

$$K_m = [S], \text{ when } V = 1/2 V_{\max}$$

This is a very useful, practical definition of K_m : K_m is equivalent to the substrate concentration at which V is one-half $V_{\max}$.

The equation describes the kinetic behavior of a great many enzymes, and all enzymes that exhibit a hyperbolic dependence of V_0 on $[S]$ are said to follow **Michaelis-Menten kinetics**.

The practical rule that $K_m = [S]$ when $V_0 = 1/2 V_{\max}$ holds for all enzymes that follow Michaelis-Menten kinetics.

The most important exceptions to Michaelis-Menten kinetics are the regulatory enzymes. 

210 Enzymes-Michaelis menten equation at low [S]

The Michaelis-Menton Equation at low [S]

Interpreting $V_{\max}$ and K_m shows a simple graphical method for obtaining an approximate value for K_m .

This graph shows the kinetic parameters that define the limits of the curve at high and low [S].

At low [S], $K_m \gg [S]$ and the $[S]$ term in the denominator of the Michaelis-Menten equation becomes insignificant.

low [S]; $K_m \gg [S]$

$$V = V_{\max} \frac{[S]}{K_m + [S]}$$

$$K_m + [S]$$

$$V = \frac{V_{\max} [S]}{K_m}$$

$$K_m$$

Since $V_{\max}$ and K_m are both constants, their ratio is a constant.

In other words, when [S] is considerably below K_m ,

V_o is proportionate to $k[S]$.

The initial reaction velocity, V_o , therefore is directly proportionate to [S].

V_o exhibits a linear dependence on [S], as observed here.

(First order Reaction)

When $[S]$ is much less than K_m (point A in Figure)

Therefore at concentrations below K_m reaction rate is first order i.e. it is directly proportional to the concentration of the substrate.

211 Enzymes-Michaelis Menten equation at high S

The Michaelis-Menton Equation at high $[S]$

At high $[S]$ $[S] \gg K_m$

The term $K_m + [S]$ is essentially equal to $[S]$.

The K_m term in the denominator of the Michaelis- Menten equation becomes insignificant

Replacing $K_m + [S]$ with $[S]$ reduces equation

high $[S]$ $[S] \gg K_m$

$$V_0 = \frac{V_{\max} [S]}{K_m + [S]}$$

$$K_m + [S]$$

Ignoring K_m

$$V_0 = \frac{V_{\max} [S]}{[S]}$$

$$[S]$$

$$V_0 = V_{\max}.$$

This is consistent with the plateau observed at high $[S]$. (Zero Order Reaction)

The rate of reaction is then independent of substrate concentration $[S]$, and is said to be **zero order** with respect to substrate concentration.

The Michaelis-Menten equation is therefore consistent with the observed dependence of V_0 on $[S]$, and the shape of the curve is defined by the terms;

V_{max}/K_m at low $[S]$ and

V_{max} at high $[S]$.

212 Enzymes-Order of reaction

Order of Reaction

When $[S]$ is much less, then the velocity of the reaction is approximately proportional to the substrate concentration.

The rate of reaction is then said to be first order with respect to substrate.

When $[S]$ is much greater than K_m the velocity is constant and equal to V_{max} .

The rate of reaction is then independent of substrate concentration, and is said to be zero order with respect to substrate concentration

Reaction Orders with Respect to Substrate Concentration

Relationship of velocity to enzyme concentration

The rate of the reaction is directly proportional to the enzyme concentration.

There is a linear relationship between reaction rate and enzyme concentration (at constant substrate concentration)

For example,

if the enzyme concentration is halved, the initial rate of the reaction (V) as well as that of V_{max} are reduced to one half that of the original.

213 Enzymes-Line Weaver Burke plot

Lineweaver-Burke plot

The Michaelis-Menten equation can be algebraically transformed into **LineweaverBurke plot**, a *Double Reciprocal Plot*, that is useful in the practical determination of K_m and V_{max} .

However, if $1/V$ is plotted versus $1/[S]$, a straight line is obtained. This graph, the LineweaverBurke plot can be used to calculate

K_m and V_{max} , as well as to determine the mechanism of action of enzyme inhibitors.

$$V = \frac{V_{max} [S]}{K_m + [S]}$$

Lineweaver-Burke transformation is derived simply by taking the reciprocal of both sides of the Michaelis-Menten equation:

$$\frac{1}{V} = \frac{K_m + [S]}{V_{max} [S]}$$

$$\frac{1}{V} = \frac{K_m}{V_{max} [S]} + \frac{1}{V_{max}}$$

$$\frac{1}{V} = \frac{K_m}{V_{max} [S]} + \frac{1}{V_{max}}$$

$$\frac{1}{V} = \frac{K_m}{V_{max} [S]} + \frac{1}{V_{max}}$$

$$\frac{1}{V} = \frac{K_m}{V_{max} [S]} + \frac{1}{V_{max}}$$

In lineweaver Burke plot y- intercept is $1/V_{\max}$ and x-intercept is $1/K_m$.

In lineweaver Burke plot y- Axis intercept is $1/V_{\max}$ and x- Axis intercept is $-1/K_m$.

Lineweaver-Burk plot, has the great advantage of allowing a more accurate determination of $V_{\max}$,

Which can only be approximated from a simple plot of V versus $[S]$.

The double-reciprocal plot of enzyme reaction rates is very useful in distinguishing between certain types of enzymatic reaction mechanisms and in analyzing enzyme inhibition.

214 Enzymes-Inhibition of Enzyme Activity

INHIBITION OF ENZYME ACTIVITY

Enzyme inhibitors are molecular agents that interfere with catalysis, slowing or halting enzymatic reactions.

Any substance that can diminish the velocity of an enzyme-catalyzed reaction is called an inhibitor.

Enzymes catalyze virtually all cellular processes, so it should not be surprising that enzyme inhibitors are among the most important pharmaceutical agents known. 

For example, aspirin inhibits the enzyme (cyclooxygenase) that catalyzes the first step in the synthesis of prostaglandins, compounds involved in many processes, including some that produce pain.

Inhibitors can be classified on the basis of their site of action on the enzyme, on whether they chemically modify the enzyme, or on the kinetic parameters they influence.

Two broad classes of enzyme inhibitors:

1. Reversible

2. Irreversible.

In general, irreversible inhibitors bind to enzymes through covalent bonds.

Reversible inhibitors typically bind to enzymes through noncovalent bonds,

The two most commonly encountered types of reversible inhibition are;

competitive and

noncompetitive.

Competitive inhibitors resemble the substrate and compete for binding to the active site of the enzyme.

Noncompetitive inhibitors do not bind at the active site. They bind either free enzyme at a site other than active site or the ES complex.

215 Enzymes-Competitive Inhibition

Competitive Inhibition

This type of inhibition occurs when the inhibitor binds reversibly to the same site that the substrate would normally occupy i.e. active site and, therefore, competes with the substrate for that site.

J ywzo'śs/o"xr \$ śy°•"l xn"y"r o"ox" –w o •"knis/o"•de.

While the inhibitor (I) occupies the active site it prevents binding of the substrate to the enzyme. Many competitive inhibitors are compounds that resemble the substrate and combine with the enzyme to form an EI complex, but without leading to catalysis.


Reversible inhibitors bind to enzymes through non covalent bonds.

Dilution of the enzyme-inhibitor complex results in dissociation of the reversibly bound inhibitor, and recovery of enzyme activity.

1. Effect on Vmax

The effect of a competitive inhibitor is reversed by increasing [S].

At a sufficiently high substrate concentration, the reaction velocity reaches the V_{max} as observed in the absence of inhibitor.



$$V_{max,app} = V_{max} \quad K_{m,app} > K_m$$

Even if high conc. of substrate is required V_{max} is achieved. $K_{m,app}$ becomes greater than K_m

When [S] far exceeds [I], the probability that an inhibitor molecule will bind to the enzyme is minimized and the reaction exhibits a normal V_{max} .


Therefore, a competitive inhibitor does not decrease V_{max}

216 Enzymes-Competitive inhibition (continue...)

2. Effect on Km

The [S] at which $V_0 = 1/2 V_{max}$, the apparent K_m , increases in the presence of inhibitor.

A competitive inhibitor increases the apparent $K_m - K_m$ for a given substrate.



$$V_{\max,app} = V_{\max} \quad K_{m,app} > K_m$$

Increase in the apparent K_m - K_m) for a given substrate means that, more substrate is now needed to achieve $1/2V_{\max}$.

3. Effect on Lineweaver-Burk plot

Competitive inhibition shows a characteristic Lineweaver-Burke plot


the plots of the inhibited and uninhibited reactions intersect at a single point on the y-axis at $1/V_{\max}$.

$V_{\max}$ is unchanged

However, the inhibited and uninhibited reactions show; different x axis intercepts

Therefore indicating that the apparent K_m is increased in the presence of the competitive inhibitor because $-1/K_m$ moves closer to zero from a negative value.

The Lineweaver-Burk plot is diagnostic for competitive inhibition



Increasing [I] results in a family of lines with a common intercept on the $1/V_0$ axis but with different slopes. Because the intercept on the $1/V_0$ axis equals $1/V_{\max}$.

Competitive inhibition.

We know that V_{max} is unchanged by the presence of a competitive inhibitor.

This is because more substrate means more formation of ES complex vs EI complex.

217 Enzymes-Non competitive inhibition

Noncompetitive Inhibition

Inhibitors bind enzymes at sites distinct from the substrate-binding site and generally bear little or no structural resemblance to the substrate.

Noncompetitive inhibitors bind at a separate site, but may bind to either E or ES.

Noncompetitive inhibition cannot be overcome by increasing the concentration of substrate.

1. Effect on V_{max}

the apparent V_{max} changes, because the inhibitor is capable of preventing catalysis regardless of whether the substrate is bound to the enzyme.

Noncompetitive inhibition cannot be overcome by increasing the concentration of substrate.

Thus, noncompetitive inhibitors decrease the V_{max} of the reaction.

V_{max} decreases in noncompetitive inhibition.

V_{max} decreases in noncompetitive inhibition.

$V_{\max}$ decreases in noncompetitive inhibition.

The noncompetitive inhibitor, in effect, lowers the concentration of the active enzyme and therefore decreases the $V_{\max}$ of the enzyme.

218 Enzymes-Non competitive inhibition (continued...)

Noncompetitive Inhibition

(Contd.)

2. Effect on K_m

Noncompetitive inhibitors do not interfere with the binding of substrate to enzyme.

Thus, the enzyme shows the same K_m in the presence or absence of the noncompetitive inhibitor.

$V_{\max}$ changes but the $[S]$ required to achieve $\frac{1}{2} V_{\max}$ i.e. K_m remains the same


3. Effect on Lineweaver-Burk plot

Noncompetitive inhibition shows a characteristic Lineweaver-Burke plot

the plots of the inhibited and uninhibited reactions intersect at a single point on the x-axis at K_m

K_m is unchanged

However, the inhibited and uninhibited reactions show; different y axis intercepts indicating the decrease in $V_{\max}$ in the presence of the competitive inhibitor.



While certain inhibitors exhibit characteristics of a mixture of competitive and noncompetitive inhibition, the evaluation of these inhibitors exceeds the scope of our level of study.

219 Water, PH and Buffers

Water

Water is the most abundant substance in living systems, making up 70% or more of the weight of most organisms

The hydrogen bonding between water molecules and the slight tendency of water to ionize are of crucial importance to the structure and function of biomolecules

Hydrogen Bonding gives Water its Unusual Properties

Water has a higher

melting point,

boiling point, and

heat of vaporization than most other common solvents

Each hydrogen atom of a water molecule shares an electron pair with the central oxygen atom


The geometry of the molecule is dictated by the shapes of the outer electron orbitals of the oxygen atom

These orbitals describe a rough tetrahedron, with a hydrogen atom at each of two corners and unshared electron pairs at the other two corners.

The oxygen nucleus attracts electrons more strongly than does the hydrogen nucleus. That is, oxygen is more electronegative.

This means that the shared electrons are more often in the vicinity of the oxygen atom than of the hydrogen

The result of this unequal electron sharing is two electric dipoles in the water molecule



the oxygen atom bears a partial negative charge equal to the sum of the two partial
charges ($\delta^- = 2 \times \delta^+$)

As a result, there is an electrostatic attraction between the oxygen atom of one water molecule and the hydrogen of another, called a hydrogen bond

The nearly tetrahedral arrangement of the orbitals about the oxygen atom allows each water molecule to form hydrogen bonds with as many as four neighboring water molecules

Each water molecule can form two hydrogen bonds involving their hydrogen atoms plus two further hydrogen bonds utilizing the hydrogen atoms attached to neighboring water molecules.

These four hydrogen bonds optimally arrange themselves tetrahedrally around each water molecule


220 Water, PH and buffer systems-ionization of water

Ionization of Water

Although many of the solvent properties of water can be explained in terms of the uncharged H₂O molecule, the small degree of ionization of water into hydrogen ions (H⁺) and hydroxide ions (OH⁻) must also be taken into account

Pure Water Is Slightly Ionized

Water molecules have a slight tendency to undergo reversible ionization to yield a hydrogen ion (a proton) and a hydroxide ion, giving the equilibrium reaction



Hydrogen ions formed in water are immediately hydrated to hydronium ions (H₃O⁺)

The ionization of water can be measured by its electrical conductivity

Pure water carries electrical current as H₃O⁺ migrates toward the cathode and OH⁻ toward the anode

The Ionization Of Water Is Expressed By an Equilibrium Constant

The degree of ionization of water at equilibrium is small

At 25°C only about two out of every 10⁹ molecules in pure water are ionized at any instant

The equilibrium constant for the reversible ionization of water is

$$K_{eq} = \frac{[\text{H}^+][\text{OH}^-]}{[\text{H}_2\text{O}]}$$

$$K_{eq} = \frac{[\text{H}^+][\text{OH}^-]}{[\text{H}_2\text{O}]}$$

Since 1 mole (mol) of water weighs 18 g,

1 liter (L) (1000 g) of water contains

$$1000 \div 18 = 55.56 \text{ mol}$$

Pure water thus is 55.56 molar(M)

Accordingly, we can substitute 55.5 M in the equilibrium constant expression to yield

$$K_{eq} = \frac{[\text{H}^+][\text{OH}^-]}{55.5}$$

$$K_{eq} = \frac{[\text{H}^+][\text{OH}^-]}{55.5}$$

On rearranging, this becomes (K_{eq}) (55.5) = [H⁺][OH⁻] = K_w

Where K_w designates the product (55.5M)(K_{eq}), the ion product of water at 25°C

The value for K_{eq} , determined by electrical conductivity measurements of pure water, is
 $1.8 \times 10^{-16} \text{ M}$ at 25°C

$$K_w = (K_{eq})(55.5) = [\text{H}^+][\text{OH}^-]$$

$$K_w = (1.8 \times 10^{-16})(55.5) = [\text{H}^+][\text{OH}^-]$$

$$K_w = 1.0 \times 10^{-14} = [\text{H}^+][\text{OH}^-]$$

$$K_w = [\text{H}^+][\text{OH}^-] = [\text{H}^+]^2 = [\text{OH}^-]^2$$

Solving for $[\text{H}^+]$ gives:

$$[\text{H}^+] = \sqrt{K_w} = 10^{-7} \text{ M}$$

$$[\text{H}^+] = 10^{-7} \text{ M}$$

Thus the product $[\text{H}^+][\text{OH}^-]$ in aqueous solutions at 25°C always equals $1 \times 10^{-14} \text{ M}^2$

When there are exactly equal concentrations of H^+ and OH^- , as in pure water, the solution is said to be at neutral pH

As the ion product of water is constant, whenever $[\text{H}^+]$ is greater than $1 \times 10^{-7} \text{ M}$, $[\text{OH}^-]$ must be less than $1 \times 10^{-7} \text{ M}$, and vice versa

221 Water, PH and Buffer systems-working examples

WORKING EXAMPLES

From the ion product of water we can calculate $[\text{H}^+]$

if we know $[\text{OH}^-]$, and vice versa

What is the concentration of H^+ in a solution of 0.1 M NaOH?

Solution:

$$K_w = [\text{H}^+][\text{OH}^-]$$

With $[\text{OH}^-] = 0.1 \text{ M}$, solving for $[\text{H}^+]$ gives

$$[\text{H}^+] = K_w / [\text{OH}^-]$$

$$= 1 \times 10^{-14} \text{ M}^2 / 0.1 \text{ M}$$

$$= 10^{-14} \text{ M}^2 / 0.1 \text{ M}$$

$$= 10^{-13} \text{ M}$$

What is the concentration of OH^- in a solution with an H^+ concentration of $1.3 \times 10^{-4} \text{ M}$?

Solution:

$$K_w = [\text{H}^+][\text{OH}^-]$$

With $[\text{H}^+] = 1.3 \times 10^{-4} \text{ M}$, solving for $[\text{OH}^-]$ gives

$$[\text{OH}^-] = K_w / [\text{H}^+]$$

$$= 1 \times 10^{-14} \text{ M}^2 / 1.3 \times 10^{-4} \text{ M}$$

$$= 7.7 \times 10^{-11} \text{ M}$$

The pH Scale
 Designates the H^+ and OH^- Concentrations

The **pH** of a solution is defined as the logarithm to the base 10 of the reciprocal of the $[\text{H}^+]$, i. e, the negative logarithm of the $[\text{H}^+]$

$$\text{pH} = \log 1/[\text{H}^+]$$

$$= -\log[\text{H}^+]$$

The pH of water at 25°C, in which H^+ and OH^- ions are present in equal numbers, is 7.0

$$\text{pH} = -\log [1 \times 10^{-7}] = 7$$

The symbol p denotes "negative logarithm of"

For each pH unit less than 7.0, the $[\text{H}^+]$ is increased tenfold;

for each pH unit above 7.0, it is decreased tenfold

What will be the pH of 0.1 M HCl?

A 0.1 M HCl solution will contain 0.1 or 10^{-1} grams H^+ per litre

$$\text{pH} = -\log [\text{H}^+]$$

$$\text{pH} = -\log [10^{-1}]$$

$$= -[-1]$$

$$= 1$$

222 Water, PH and buffer systems-weak acids and bases

Weak Acids and Bases

Each acid has a characteristic tendency to ionize in an aqueous solution

The stronger the acid, the greater its tendency ionize i.e. to lose its proton

This tendency is measured by an acid dissociation constant

Weak Acids and Bases have Characteristic Acid Dissociation Constants

HCl, H₂SO₄, and HNO₃, commonly called strong acids, are fully ionized in aqueous solutions

The strong bases NaOH and KOH are also completely ionized


Of more interest is the behavior of weak acids and bases-those not completely ionized when dissolved in water

These are ubiquitous in biological systems and play important roles in metabolism and its regulation


Acids may be defined as proton donors and bases as proton acceptors

A proton donor and its corresponding proton acceptor make up a conjugate acidbase pair

Acetic acid (CH₃COOH), a proton donor, and the acetate anion (CH₃COO⁻), the corresponding proton acceptor, constitute a conjugate acid-base pair, related by the reversible reaction:



The tendency of any acid (HA) to lose a proton and form its conjugate base (A⁻) is defined by the acid dissociation constant (K_a) for the reversible reaction



$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

Stronger acids, have larger dissociation constants (K_a) i.e they ionize completely

Weaker acids, have smaller dissociation constants (K_a) i.e. they ionize only partially.

pK_a

analogous to pH, pK_a is defined by the equation

$$\text{pK}_a = \log 1/K_a$$

$$= -\log K_a$$

The stronger the tendency to dissociate a proton, the stronger is the acid and the lower its pK_a

223 Water, PH and buffer systems-working with pK_a

Working with pK_a

Titration is used to determine the amount of an acid in a given solution

A measured volume of the acid is titrated with a solution of a strong base, usually sodium hydroxide (NaOH), of known concentration

Consider the titration of a 0.1M solution of acetic acid with 0.1M NaOH at 25°C

Two reversible equilibria are involved in the process (here, for simplicity, acetic acid is denoted HAc) o H₂O ⇌ H⁺ + OH⁻



The equilibria must simultaneously conform to their characteristic equilibrium constants, which are, respectively,

$$K_w = [H^+][OH^-] = 1 \times 10^{-14} \text{ M}^2 \quad K_a = \frac{[H^+][Ac^-]}{[HAc]} = 1.74 \times 10^{-5} \text{ M}$$

At the beginning of the titration, the acetic acid is only slightly ionized

As NaOH is gradually added, OH^- combines with the free H^+ in the solution to form

H_2O As free H^+ is removed, HAc dissociates further to satisfy its equilibrium constant

The net result as the titration proceeds is that more and more HAc ionizes, forming Ac^- , as the NaOH is added

At the midpoint of the titration, at which exactly 0.5 equivalent of NaOH has been added, onehalf of the original acetic acid has undergone dissociation, so that the concentration of the proton donor, [HAc], now equals that of the proton acceptor, [Ac^-]

At the midpoint a very important relationship holds:

The pH of the equimolar solution of acetic acid and acetate is exactly equal to the pK_a of acetic acid ($pK_a = 4.76$)

At pK_a of a weak acid, half of the acid is in dissociated form whereas other half is un-dissociated.

Increasing the pH will result in an increased dissociation and vice versa.

224 Water, PH and buffer systems-HH equation

The Henderson-Hasselbalch (HH) Equation

The HH equation relates pH, pK_a , and buffer concentration

This equation is simply a useful way of restating the expression for the ionization constant of an acid

For the ionization of a weak acid HA, the HH equation can be derived as follows:

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

First solve for $[H^+]$:

$$[H^+] = K_a \frac{[HA]}{[A^-]}$$

Then take the negative logarithm of both sides:

$$-\log[H^+] = -\log K_a - \log \frac{[HA]}{[A^-]}$$

$$\text{Substitute pH for } -\log [H^+] \text{ and } pK_a \text{ for } -\log K_a \quad \text{pH} = pK_a + \log \frac{[HA]}{[A^-]}$$

Now invert $-\log [HA]/[A^-]$,

$$\text{pH} = \text{pKa} + \log[\text{A}^-]/[\text{HA}]$$

This equation shows why the pKa of a weak acid is equal to the pH of the solution at the midpoint of its titration

At that point, $[\text{HA}] = [\text{A}^-]$

$$\text{pH} = \text{pKa} + \log[\text{A}^-]/[\text{HA}]$$

$$\text{pH} = \text{pKa} + \log 1$$

$$\text{pH} = \text{pKa} + 0$$

$$\text{pH} = \text{pKa}$$

Calculate the pKa of lactic acid, given that when the concentration of lactic acid is 0.01M and the concentration of lactate is 0.087 M, the pH is 4.80

$$\text{pH} = \text{pKa} + \log [\text{lactate}]/[\text{lactic acid}]$$

$$\text{pKa} = \text{pH} - \log [\text{lactate}]/[\text{lactic acid}]$$

$$= 4.80 - \log 0.087/0.01$$

$$= 4.80 - \log 8.7$$

$$= 4.80 - 0.94$$

$$= \mathbf{3.9}$$

Calculate the ratio of the concentrations of acetate and acetic acid required in a buffer system of pH 5.30 (pKa = 4.76)

$$\log [\text{acetate}]/[\text{acetic acid}] = \text{pH} - \text{pKa}$$

$$= 5.30 - 4.76$$

$$= 0.54$$

$$[\text{acetate}]/[\text{acetic acid}] = \text{antilog } 0.54$$

$$= \mathbf{3.5}$$

In summary,

when $[\text{HA}] = [\text{A}^-]$; $\text{pH} = \text{pKa}$

when $[\text{HA}] > [\text{A}^-]$; $\text{pH} < \text{pKa}$

when $[\text{HA}] < [\text{A}^-]$; $\text{pH} > \text{pKa}$

225 Water, PH and buffer systems-buffer solutions

Buffer Solutions

Buffers are aqueous systems that tend to resist changes in pH when small amounts of acid (H^+) or base (OH^-) are added

A buffer system consists of a weak acid (the proton donor) and its conjugate base (the proton acceptor)

As an example, a mixture of acetic acid and acetate ion, is a buffer system,

If more H^+ is added to this solution, it simply shifts the equilibrium to the left, absorbing H^+ , so the $[\text{H}^+]$ remains unchanged.

If H^+ is removed (e.g. by adding OH^-) then the equilibrium shifts to the right, releasing H^+ to keep the pH constant

Notice that the titration curve of acetic acid has a relatively flat zone

extending about 1 pH unit on either side of its midpoint pH of 4.76.

In this zone, a given amount of H^+ or OH^- added to the system has much less effect on pH than the same amount added outside the zone

This relatively flat zone is the buffering region of the acetic acid-acetate buffer pair

At the midpoint of the buffering region, where the concentration of the proton donor (acetic acid) exactly equals that of the proton acceptor (acetate), the buffering power is maximal;

The pH at this point in the titration curve of acetic acid is equal to its pK_a

The pH of the acetate buffer system does change slightly when a small amount of H^+ or OH^- is added, but this pH change is very small

compared with the pH change that would result if the same amount of H^+ or OH^- were added to pure water.

Each conjugate acid-base pair has a characteristic pH zone in which it is an effective buffer

For example, the $\text{H}_2\text{PO}_4^-/\text{HPO}_4^{2-}$ pair has a pK_a of 6.86 and thus can serve as an effective buffer system between approximately pH 5.9 and pH 7.9

